



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12N 9/99, 9/50, A01N 43/02, 43/06, 43/08, 43/16, 43/18, C07D 307/00, 307/48, 309/00, 319/06, 321/00, 333/02, 333/12, 333/22, 335/00, 335/02</p>	A1	<p>(11) International Publication Number: WO 98/50533</p> <p>(43) International Publication Date: 12 November 1998 (12.11.98)</p>
<p>(21) International Application Number: PCT/US98/03200</p> <p>(22) International Filing Date: 6 May 1998 (06.05.98)</p> <p>(30) Priority Data: 60/045,758 6 May 1997 (06.05.97) US</p> <p>(71) Applicants (for all designated States except US): SMITHK- LINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). SMITHK- LINE BEECHAM PLC [GB/GB]; Great West Road, New Horizons Court, Brentford, Middlesex TW8 9BD (GB).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): GRIBBLE, Andrew, D. [GB/GB]; 1 Wolvescroft, Woolmer Green Knebworth, Herts SG3 6LR (GB). FENWICK, Ashley, Edward [GB/GB]; 27 Whitecroft Road, Meldreth, Cambridgeshire SG8 6NG (GB). MARQUIS, Robert, W. [US/US]; 115 Cambria Court, St. Davids, PA 19087 (US). VEBER, Daniel, F. [US/US]; 290 Batleson Road, Ambler, PA 19002 (US). WITHER- INGTON, Jason [GB/GB]; 32 West Road, Sawbridgeworth, Hertfordshire CM21 0BL (GB).</p>	<p>(74) Agents: McCARTHY, Mary, E. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).</p> <p>(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: PROTEASE INHIBITORS</p> <p>(57) Abstract</p> <p>The invention relates to 3-hydroxy- and 3-keto-cyclohetero-substituted leucine compounds that are inhibitors of cysteine proteases, particularly cathepsin K, and are useful in the treatment of diseases in which inhibition of bone loss is a factor. The 3-hydroxy- or 3-keto-moiety is bonded to a tetrahydrothiophene, tetrahydrothiopyran, tetrahydrofuran or tetrahydropyran ring.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROTEASE INHIBITORS

Field of the Invention

This invention relates to novel protease inhibitors, particularly inhibitors of
5 cysteine and serine proteases, more particularly compounds which inhibit cysteine
proteases. The compounds of this invention even more particularly relate to those
compounds which inhibit cysteine proteases of the papain superfamily, and particularly
cysteine proteases of the cathepsin family. In the most preferred embodiment, this
invention relates to compounds which inhibit cathepsin K. Such compounds are
10 particularly useful for treating diseases in which cysteine proteases are implicated,
especially diseases of excessive bone or cartilage loss, e.g., osteoporosis, periodontitis,
and arthritis.

Background of the Invention

15 Cathepsin K is a member of the family of enzymes which are part of the papain
superfamily of cysteine proteases. Cathepsins B, H, L, N and S have been described in
the literature. Recently, cathepsin K polypeptide and the cDNA encoding such
polypeptide were disclosed in U.S. Patent No. 5,501,969 (called cathepsin O therein).
Cathepsin K has been recently expressed, purified, and characterized. Bossard, M. J., et
20 al., (1996) *J. Biol. Chem.* **271**, 12517-12524; Drake, F.H., et al., (1996) *J. Biol. Chem.*
271, 12511-12516; Bromme, D., et al., (1996) *J. Biol. Chem.* **271**, 2126-2132.

Cathepsin K has been variously denoted as cathepsin O, cathepsin X or
cathepsin O2 in the literature. The designation cathepsin K is considered to be the more
appropriate one (name assigned by Nomenclature Committee of the International Union
25 of Biochemistry and Molecular Biology).

Cathepsins of the papain superfamily of cysteine proteases function in the
normal physiological process of protein degradation in animals, including humans, e.g.,
in the degradation of connective tissue. However, elevated levels of these enzymes in
the body can result in pathological conditions leading to disease. Thus, cathepsins have
30 been implicated in various disease states, including but not limited to, infections by
pneumocystis carinii, *trypanoma cruzi*, *trypanoma brucei*, and *Crithidia*
fusculata; as well as in schistosomiasis malaria, tumor metastasis, metachromatic
leukodystrophy, muscular dystrophy, amyotrophy, and the like. See International
Publication Number WO 94/04172, published on March 3, 1994, and references cited
35 therein. See also European Patent Application EP 0 603 873 A1, and references cited
therein. Two bacterial cysteine proteases from *P. gingivalis*, called gingipains, have

been implicated in the pathogenesis of gingivitis. Potempa, J., et al. (1994) *Perspectives in Drug Discovery and Design*, **2**, 445-458.

5 Cathepsin K is believed to play a causative role in diseases of excessive bone or cartilage loss. Bone is composed of a protein matrix in which spindle- or plate-shaped crystals of hydroxyapatite are incorporated. Type I Collagen represents the major structural protein of bone comprising approximately 90% of the structural protein. The remaining 10% of matrix is composed of a number of non-collagenous proteins, including osteocalcin, proteoglycans, osteopontin, osteonectin, thrombospondin, fibronectin, and bone sialoprotein. Skeletal bone undergoes remodeling at discrete foci throughout life. These foci, or remodeling units, undergo a cycle consisting of a bone resorption phase followed by a phase of bone replacement.

10 Bone resorption is carried out by osteoclasts, which are multinuclear cells of hematopoietic lineage. The osteoclasts adhere to the bone surface and form a tight sealing zone, followed by extensive membrane ruffling on their apical (i.e., resorbing) surface. This creates an enclosed extracellular compartment on the bone surface that is acidified by proton pumps in the ruffled membrane, and into which the osteoclast secretes proteolytic enzymes. The low pH of the compartment dissolves hydroxyapatite crystals at the bone surface, while the proteolytic enzymes digest the protein matrix. In this way, a resorption lacuna, or pit, is formed. At the end of this phase of the cycle, osteoblasts lay down a new protein matrix that is subsequently mineralized. In several disease states, such as osteoporosis and Paget's disease, the normal balance between bone resorption and formation is disrupted, and there is a net loss of bone at each cycle. Ultimately, this leads to weakening of the bone and may result in increased fracture risk with minimal trauma.

25 The abundant selective expression of cathepsin K in osteoclasts strongly suggests that this enzyme is essential for bone resorption. Thus, selective inhibition of cathepsin K may provide an effective treatment for diseases of excessive bone loss, including, but not limited to, osteoporosis, gingival diseases such as gingivitis and periodontitis, Paget's disease, hypercalcemia of malignancy, and metabolic bone disease. Cathepsin K levels have also been demonstrated to be elevated in chondroclasts of osteoarthritic synovium. Thus, selective inhibition of cathepsin K may also be useful for treating diseases of excessive cartilage or matrix degradation, including, but not limited to, osteoarthritis and rheumatoid arthritis. Metastatic neoplastic cells also typically express high levels of proteolytic enzymes that degrade the surrounding matrix. Thus, selective inhibition of cathepsin K may also be useful for treating certain neoplastic diseases.

It now has been discovered that a novel class of compounds are protease inhibitors, most particularly inhibitors of cathepsin K, and these compounds are useful for treating diseases in which inhibition of bone resorption is indicated, such as osteoporosis and periodontal disease.


5

Summary of the Invention

An object of the present invention is to provide protease inhibitors, such as inhibitors of cysteine and serine proteases. In particular, the present invention relates to compounds which inhibit cysteine proteases, and particularly cysteine proteases of the papain superfamily. Preferably, this invention relates to compounds which inhibit cysteine proteases of the cathepsin family and particularly, compounds which inhibit cathepsin K. The compounds of the present invention are useful for treating diseases, which may be therapeutically modified by altering the activity of such proteases.

10

Accordingly, in the first aspect, this invention provides a compound according to formula (I):



20

In another aspect, this invention provides a pharmaceutical composition comprising a compound according to formula (I) and a pharmaceutically acceptable carrier.


In yet another aspect, this invention provides a method of treating diseases in which the disease pathology may be therapeutically modified by inhibiting proteases, such as cysteine and serine proteases. In particular, the method includes treating diseases by inhibiting cysteine proteases, and particularly cysteine proteases of the papain superfamily. More particularly, the inhibition of cysteine proteases of the cathepsin family, such as cathepsin K is described.

In another aspect, the compounds of this invention are especially useful for treating diseases characterized by bone loss, such as osteoporosis and gingival diseases, such as gingivitis and periodontitis, or by excessive cartilage or matrix degradation, such as osteoarthritis and rheumatoid arthritis.

30

Detailed Description of the Invention

The present invention provides compounds of formula (I):



wherein:

R^1 is R'' , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

R^2 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;

R^3 is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆cycloalkyl-C₀₋₆alkyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;

R^4 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;

each R^5 independently is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;

R^6 is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl-C₀₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl;

R' is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;

R'' is C₁₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, Ar-C₂₋₆alkenyl or Het-C₂₋₆alkenyl;

X is O or S; and

n is 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

The present invention includes all hydrates, solvates, complexes and prodrugs of the compounds of this invention. Prodrugs are any covalently bonded compounds which release the active parent drug according to formula (I) *in vivo*. If a chiral center or another form of an isomeric center is present in a compound of the present invention, all forms of such isomer or isomers, including enantiomers and diastereomers, are intended to be covered herein. Inventive compounds containing a chiral center may be used as a racemic mixture, an enantiomerically enriched mixture, or the racemic mixture may be separated using well-known techniques and an individual enantiomer may be used alone. According to the instant invention, the S-form at the furan ring junction of formula (I) compounds is preferred.

- In cases in which compounds have unsaturated carbon-carbon double bonds, both the *cis* (*Z*) and *trans* (*E*) isomers are within the scope of this invention. In cases wherein compounds may exist in tautomeric forms, such as keto-enol tautomers, each tautomeric form is contemplated as being included within this invention whether existing in equilibrium or predominantly in one form.

The meaning of any substituent at any one occurrence in formula (I) or any subformula thereof is independent of its meaning, or any other substituent's meaning, at any other occurrence, unless specified otherwise.

With respect to formula (I):


10

Suitably, R^2 and R^4 are H and R^3 is C_{1-6} alkyl or C_{2-6} alkenyl. Preferably, R^3 is *i*-butyl.

Suitably, each R^5 is H.

Suitably, R^1 is $R''OC(O)$, $R''SO_2$ or $R''C(O)$, in which R'' is $Ar-C_{0-6}$ alkyl or


- 15 Het- C_{0-6} alkyl, and, most preferably, R'' is




in which B_2 is OH, CN, OCF_3 , OC_{1-6} alkyl, OAr, SO_2C_{1-6} alkyl, C_{1-6} alkyl or halo.

- 20 Suitably, *n* is 1 or 2. Preferably, *n* is 1.
Suitably, X is O.


In one particular embodiment, this invention is a compound of formula (II):



5 Preferably, the formula (I) compound of this invention is a compound of formula (IIa):




Alternately, the formula (I) compound of this invention is a compound of formula (IIb):



10

In another embodiment, this invention is a compound of formula (IIc):



15

Specific representative compounds of this invention are:

4-(*R,S*)-Amino-N-[(3,4-methylenedioxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;

4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;

4-(*R,S*)-Amino-N-[(3,4-dichlorobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;

4-(*R,S*)-Amino-N-[(2-quinolinecarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;


20

4-(*R,S*)-Amino-N-[(8-quinolinecarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;

4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-2,2-dibenzyl-tetrahydrofuran-3-one;

- 4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(indole-6-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 5 4-(*R,S*)-Amino-N-[(benzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(5-aminobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(5-chlorobenzo[furan-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 10 4-(*R,S*)-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(3-bromobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(4-bromobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(5-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 15 4-(*R,S*)-Amino-N-[(4-fluorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(4-phenoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(4-phenylbenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 20 4-(*R,S*)-Amino-N-[(6-trifluoromethylbenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(4-ethylbenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(4-(tert-butyl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(5-methoxybenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 25 4-(*R,S*)-Amino-N-[(4-nitrobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(6-bromobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 30 4-(*R,S*)-Amino-N-[(5-bromobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-(*R,S*)-Amino-N-[(6-methoxybenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 4-*S*-Amino-N-[(benzo(b)thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 35 one;
- 4-*R*-Amino-N-[(benzo(b)thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;

- 4-*S*-Amino-N-[(2-naphthyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*R*-Amino-N-[(2-naphthyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(quinoline-2-carbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*R*-Amino-N-[(quinoline-2-carbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 5 4-*S*-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-*S*-leucine]-
 tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-pyrid-3-yl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-pyrid-2-yl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(benzylloxycarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 10 4-*S*-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(benzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-[6-methylpyrid-3-yl]benzoyl)-*S*-leucine]-tetrahydrofuran-3-
 one;
 4-*S*-Amino-N-[(5-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-
 15 tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-pyrid-4-yl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(2-chlorobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-bromobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-
 20 tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-benzylpiperidin-1-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-
 one;
 4-*S*-Amino-N-[(3,4-dichlorobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(3-chlorobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 25 4-*(R,S)*-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-tetrahydropyran-3-one;
 4-*(R,S)*-Amino-N-[(4-phenoxybenzoyl)-*S*-leucine]-tetrahydropyran-3-one;
 4-*(R,S)*-Amino-N-[(quinolin-2-ylcarbonyl)-*S*-leucine]-tetrahydropyran-3-one;
 4-*(R,S)*-Amino-N-[(benzylloxycarbonyl)-*S*-leucine]-tetrahydropyran-3-one;
 4-*(R,S)*-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-
 30 tetrahydropyran-3-one; and
 4-*(R,S)*-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-
 tetrahydrothiophen-3-one;
 or a pharmaceutically acceptable salt thereof.
- In yet another aspect, this invention provides novel intermediates useful in the
 35 preparation of formula (I) compounds represented by formulae (III), (IV) and (V):



wherein:

R^1 is R'' , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

5 R^2 is H, C_{1-6} alkyl, C_{2-6} alkenyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

R^3 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

R^4 is H, C_{1-6} alkyl, C_{2-6} alkenyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

10 each R^5 independently is H, C_{1-6} alkyl, C_{2-6} alkenyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

R^6 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, Ar- C_{0-6} alkyl,

Het- C_{0-6} alkyl;


R' is H, C_{1-6} alkyl, C_{2-6} alkenyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

R'' is C_{1-6} alkyl, Ar- C_{0-6} alkyl, Het- C_{0-6} alkyl, Ar- C_{2-6} alkenyl; or

15 Het- C_{2-6} alkenyl; and

n is 1, 2 or 3;

or a pharmaceutically acceptable salt thereof, or



20 wherein:

R^1 is R'' , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

R^2 is H, C_{1-6} alkyl, C_{2-6} alkenyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

25 R^3 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

R^4 is H, C_{1-6} alkyl, C_{2-6} alkenyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

each R^5 independently is H, C_{1-6} alkyl, C_{2-6} alkenyl, Ar- C_{0-6} alkyl, or Het- C_{0-6} alkyl;

R^6 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, Ar- C_{0-6} alkyl,


30 Het- C_{0-6} alkyl;

R' is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;

R" is C₁₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, Ar-C₂₋₆alkenyl; or
Het-C₂₋₆alkenyl; and

n is 1, 2 or 3;

- 5 or a pharmaceutically acceptable salt thereof, or



wherein:

- R¹ is R", R" C(O), R" C(S), R" SO₂, R" OC(O), R" R' NC(O), or
10 R" OC(O) NR' CH(R⁶) C(O);
R³ is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆cycloalkyl-C₀₋₆alkyl,
Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
R⁴ is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
each R⁵ independently is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or
15 Het-C₀₋₆alkyl;
R⁶ is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl-C₀₋₆alkyl, Ar-C₀₋₆alkyl,
Het-C₀₋₆alkyl;
R' is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
R" is C₁₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, Ar-C₂₋₆alkenyl; or
20 Het-C₂₋₆alkenyl;
X is O or S; and
n is 1, 2 or 3;

or a pharmaceutically acceptable salt thereof.

Representative intermediates of this invention are:

- 25 *trans*-4-(R,S)-Amino-N-[(benzyloxycarbonyl)-S-leucine]-3-
hydroxytetrahydrofuran;
trans-4-(R,S)-Amino-N-[(*tert*-butoxycarbonyl)-S-leucine]-3-
hydroxytetrahydrofuran;
trans-4-(R,S)-Amino-N-(S-leucine)-3-hydroxytetrahydrofuran;
30 *trans*-4-(R,S)-Amino-N-[(3,4-methylenedioxybenzoyl)-S-leucine]-3-
hydroxytetrahydrofuran;
trans-4-(R,S)-Amino-N-[(3,4-dichlorobenzoyl)-S-leucine]-3-
hydroxytetrahydrofuran;

- trans*-4-(*R,S*)-Amino-N-[(2-quinolinecarbonyl)-*S*-leucine]-3-hydroxytetrahydrofuran;
- trans*-4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3-hydroxytetrahydropyran;
- 5 *trans*-4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-hydroxytetrahydropyran;
- trans*-4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-hydroxytetrahydrofuran;
- trans*-4-(*R,S*)-Amino-N-[(indole-6-ylcarbonyl)-*S*-leucine]-3-
- 10 hydroxytetrahydrofuran;
- trans*-4-(*R,S*)-Amino-N-[(5-aminobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-hydroxytetrahydrofuran
- trans*-4-Amino-3-hydroxytetrahydrofuran;
- trans*-3-Hydroxy-4-benzyloxycarbonylamino-tetrahydrofuran;
- 15 4-Benzyloxycarbonylamino-tetrahydrofuran-3-one;
- 3,3-Dimethoxy-4-benzyloxycarbonylamino-tetrahydrofuran;
- 3,3-Dimethoxy-4-amino-tetrahydrofuran
- trans*-4-*S*-Amino-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3-*R*-
- 20 hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-(*S*-leucine)-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(3,4-dichlorobenzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(2-quinolinecarbonyl)-*S*-leucine]-3-*R*-
- 25 hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(benzofuran-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- 30 *trans*-4-*S*-Amino-N-[(2-naphthoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(5-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- 35 *trans*-4-*S*-Amino-N-[(4-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(4-bromobenzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;

trans-4-*S*-Amino-N-[(4-(pyrid-2-yl)benzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;

trans-4-*S*-Amino-N-[(4-(pyrid-3-yl)benzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;

5 *trans*-4-*S*-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;

trans-4-Amino-3-hydroxytetrahydropyran;

trans-4-(*R,S*)-Amino-N-[(benzyl)oxycarbonyl]-*S*-leucine]-3-hydroxytetrahydropyran;

10 *trans*-4-(*R,S*)-Amino-N-(*S*-leucine)-3-hydroxytetrahydropyran;

trans-4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-hydroxytetrahydropyran;

N-benzo[b]thiophene-2-ylcarbonyl-L-leucine methyl ester;

N-benzo[b]thiophene-2-ylcarbonyl-L-leucine;

15 *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine-*S*-(methoxycarbonylmethyl)-L-D-cysteine ethyl ester; and


2-Methoxycarbonyl-4-(*R,S*)-Amino-N-[(benzo[b]thiophene-2-ylcarbonyl)-*S*-leucine]-tetrahydrothiophene-3-one;

or salts thereof.

20 These intermediates are prepared using methods analogous to that described in Schemes 1-4 and the Examples described hereinafter.

Prodrugs of compounds of the present invention may be a prodrug of the ketone functionality of formula (I) compounds, specifically ketals or hemiketals, of the formula (VI):

25



wherein:

30 R^1 is R'' , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

R^2 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

R^3 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- R^4 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
each R^5 independently is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or
Het-C₀₋₆alkyl;
- R^6 is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl-C₀₋₆alkyl, Ar-C₀₋₆alkyl,
5 Het-C₀₋₆alkyl;
 R^7 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
 R^8 is C₁₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, Ar-C₂₋₆alkenyl; or
Het-C₂₋₆alkenyl;
X is O or S;
- 10 n is 1, 2 or 3; and
 R^a and R^a' independently are H or C₁₋₂alkyl, with the proviso that when one of
 R^a or R^a' is H, the other is C₁₋₂alkyl; or together are (CH₂)₂₋₃ forming a 5- or
6-membered ring;
or a pharmaceutically acceptable salt thereof.
- 15 Abbreviations and symbols commonly used in the peptide and chemical arts are
used herein to describe the compounds of the present invention. In general, the amino
acid abbreviations follow the IUPAC-IUB Joint Commission on Biochemical
Nomenclature as described in *Eur. J. Biochem.*, 158, 9 (1984). The term "amino acid"
as used herein refers to the D- or L- isomers of alanine, arginine, asparagine, aspartic
20 acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine,
methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine.
"C₁₋₆ alkyl" as applied herein is meant to include substituted and unsubstituted
methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl, pentyl, n-pentyl,
isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof. Any C₁₋₆alkyl
25 group may be optionally substituted independently by one or two halogens, SR', OR',
N(R')₂, C(O)N(R')₂, carbamyl or C₁₋₄alkyl, where R' is H or C₁₋₆alkyl. C₀alkyl means
that no alkyl group is present in the moiety. Thus, Ar-C₀alkyl is equivalent to Ar.
"C₃₋₆ cycloalkyl" as applied herein is meant to include substituted (i.e., alkyl,
OR, SR or halogen) and unsubstituted cyclopropane, cyclobutane, cyclopentane, and
30 cyclohexane.
"C₂₋₆ alkenyl" as applied herein means an alkyl group of 2 to 6 carbons,
wherein a carbon-carbon single bond is replaced by a carbon-carbon double bond.
C₂₋₆alkenyl includes ethylene, 1-propene, 2-propene, 1-butene, 2-butene, isobutene and
the several isomeric pentenes and hexenes. Both cis and trans isomers are included.
35 "C₂₋₆ alkynyl" means an alkyl group of 2 to 6 carbons, wherein one carbon-
carbon single bond is replaced by a carbon-carbon triple bond. C₂₋₆ alkynyl includes

acetylene, 1-propyne, 2-propyne, 1-butyne, 2-butyne, 3-butyne, and the simple isomers of pentyne and hexyne.

"Halogen" or "halo" means F, Cl, Br, and I.

"Ar" or "aryl" means unsubstituted phenyl or naphthyl; or phenyl or naphthyl

- 5 substituted by one or more of Ph-C₀₋₆alkyl, Het-C₀₋₆alkyl, C₁₋₆alkoxy, Ph-C₀₋₆alkoxy, Het-C₀₋₆alkoxy, OH, (CH₂)₁₋₆NR'R', O(CH₂)₁₋₆NR'R'; wherein each R' independently is H, C₁₋₆alkyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl; or phenyl or naphthyl substituted by one to three moieties selected from C₁₋₄alkyl, OR', N(R')₂, SR', CF₃, NO₂, CN, CO₂R', CON(R')₂, F, Cl, Br and I, or substituted by a
- 10 methylenedioxy group.

- As used herein "Het" or "heterocyclic" represents a stable 5- to 7-membered monocyclic or a stable 7- to 10-membered bicyclic heterocyclic ring, which is either saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen
- 15 and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure, and may optionally be substituted with one or two moieties selected from
- 20 C₁₋₄alkyl, OR', N(R')₂, SR', CF₃, NO₂, CN, CO₂R', CON(R')₂, F, Cl, Br and I, where R' is as defined herein before. Examples of such heterocycles include piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, thienyl, pyrrolyl, 4-piperidinyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, pyridyl, pyrazinyl, oxazolidinyl, oxazoliny, oxazolyl, isoxazolyl,
- 25 morpholinyl, thiazolidinyl, thiazolinyl, isothiazolyl, thiazolyl, quinuclidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzothienyl, benzopyranyl, benzoxazolyl, benzofuranyl, furyl, pyranyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzoxazolyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, oxadiazolyl, benzothiazolyl, benzoisothiazolyl, benzisoxazolyl, pyrimidinyl, cinnolinyl, quinazolinyl, quinoxalinyl,
- 30 1,5-naphthyridinyl, 1,6-naphthyridinyl, 1,7-naphthyridinyl, 1,8-naphthyridinyl, tetrazolyl, 1,2,3-triazolyl, and 1,2,4-triazolyl.


- Certain radical groups are abbreviated herein. t-Bu refers to the tertiary butyl radical; Boc or BOC refers to the t-butyloxycarbonyl radical; Fmoc refers to the fluorenylmethoxycarbonyl radical; Ph refers to the phenyl radical; and Cbz or CBZ
- 35 refers to the benzyloxycarbonyl radical.

Certain reagents are abbreviated herein. DCC refers to dicyclohexylcarbodiimide; EDC or EDCI refers to N-ethyl-N'-(dimethylaminopropyl)-

carbodiimide. HOBt or HOBt refers to 1-hydroxybenzotriazole; DMF refers to dimethyl formamide; DIEA refers to di-isopropylethylamine; Lawesson's reagent is 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide; TFA refers to trifluoroacetic acid; and THF refers to tetrahydrofuran.

5 Compounds of the formula (I) are generally prepared by:

(i) reacting a compound of the formula (III):




(III)

10

wherein R^1 , R^2 , R^3 , R^4 , R^5 and n are as defined in formula (I), with any reactive functional groups protected, with an oxidizing agent; or

(ii) decarboxylating a compound of the formula (IV):

15




(IV)

wherein R^1 , R^2 , R^3 , R^4 , R^5 and n are as defined in formula (I), with any reactive functional groups protected; or

20

(iii) reacting a compound of the formula (V):



25


(V)

wherein R^1 , R^3 , R^4 , R^5 and n are as defined in formula (I), with any reactive functional groups protected, with an acid;

- and thereafter removing any protecting groups and optionally forming a
5 pharmaceutically acceptable salt.

Compounds of the formula (I) are prepared by methods analogous to those described in the solution synthesis method of Scheme 1, or the solid support method of Scheme 2, or the solution synthesis method of Scheme 3.

Scheme 1




- 5 a) NaN_3 , ammonium chloride, methanol:water; b) 10% Pd/C, EtOH, H_2 ; ethanolic HCl; c) trimethylacetyl chloride, N-BOC-leucine, DIEA, CH_2Cl_2 ; d) TFA, CH_2Cl_2 ; e) RCOCl, sodium hydrogen carbonate, 1,4-dioxane; f) Dess-Martin periodinane, CH_2Cl_2


Compounds of the general formula (I), wherein n is 1 and R^1 is $\text{R}^n\text{C}(\text{O})$ are prepared by methods shown in Scheme 1. Treatment of the known epoxide 1-Scheme-1 with sodium azide and ammonium chloride in aqueous methanol at elevated temperatures provides the azide 2-Scheme-1. Reduction of the azide 2-Scheme-1 utilizing methods that are known in the art, such as reduction with palladium on carbon in ethanol under an atmosphere of hydrogen, provides the amine salt 3-Scheme-1 after treatment with ethanolic hydrogen chloride. The amine salt 3-Scheme-1 may be coupled with a carboxylic acid by methods that are known in the art, such as acylation with an acid chloride or coupling with an acid in the presence of EDC and HOBT, to provide the amide 4-Scheme-1. The *tert*-butoxycarbonyl group may be removed by treatment with a strong acid, such as TFA, in an aprotic solvent, such as

dichloromethane, to provide 5-Scheme-1. The salt 5-Scheme-1 may be acylated with an acid chloride in 1,4-dioxane in the presence of an aqueous base, such as saturated sodium hydrogen carbonate, to yield 6-Scheme-1. The alcohol 6-Scheme-1 may be oxidized by methods known in the art, such as by treatment with Dess-Martin periodinane, in an aprotic solvent, such as dichloromethane.

Scheme 2




Scheme 2 (cont.)



Compounds of the general formula (I), wherein R^1 is $C(O)R'$, R^2 is H, R^4 is H, R^5 is H, X is O, and n is 1, are prepared by the solid support synthesis (SPS) method shown in Scheme 2. In particular, in the first step, sodium triacetoxyborohydride is added to a stirred solution of Ellmans resin in DMF containing 1% HOAc, and then the α -amino acid, methyl ester is added to give rise to 2-Scheme-2. The amine group is coupled with a carboxylic acid by using known methods, such as by the addition of the carboxylic acid with EDC, to give rise to the amide, 3-Scheme-2. Thereafter, the ester group is hydrolysed, for example, with potassium trimethylsilanoate in THF, and the liberated acid group is coupled with 3,3-dimethoxy-4-amino-tetrahydrofuran using, for example, EDC in NMP, to give rise to 4-Scheme-2. The blocking group is then removed by known cleaving methods, such as by the addition of 7:2:1 TFA/ CH_2Cl_2 / H_2O , to give rise to the desired compound, 5-Scheme-2.

Compounds of the general formula (I), wherein R^1 is $C(O)R'$, R^2 is H, R^4 is H, R^5 is H, X is O, and n is 2, are prepared by the solid support synthesis (SPS) method shown in Scheme 2, except 3,3-dimethoxy-4-aminotetrahydropyran is substituted for 3,3-dimethoxy-4-amino-tetrahydrofuran.


Scheme 3



(a) L-Leu methyl ester, Pr_2NEt , CH_2Cl_2 ; (b) LiOH , $\text{THF}/\text{H}_2\text{O}$; (c) (i) ClCO_2Pr , Et_3N , CH_2Cl_2 (ii) L-Cys ethyl ester (iii) $\text{BrCH}_2\text{CO}_2\text{Me}$; (d) (i) NaOMe , MeOH , (ii) AcOH/HCl , H_2O

Scheme 4, hereinafter, shows the preparation of certain formula (I) compounds which are S-diastereomers at the furan ring junction.

Scheme 4




The R-diastereomer is prepared in an identical way from the opposite 3-azido-4-hydroxytetrahydrofuran enantiomer:



The intermediates of the present invention, such as 3,3-dimethoxy-4-aminotetrahydrofuran, can be prepared according to the method of Scheme 5.

Scheme 5



The steps in Scheme 5 are presented as follows. First, a nitrogen-protecting group is added by reacting 3-hydroxy-4-aminotetrahydrofuran with benzylchloroformate in dioxane containing aqueous sodium carbonate. Thereafter, the alcohol group is oxidized by known methods, such as by the addition of bleach containing sodium bicarbonate in the presence of sodium bromide and TEMPO in EtOAc, toluene, and water, to give rise to the ketone. The ketone is converted to the dimethylketal by the addition of trimethyl-orthoformate in the presence of paratoluenesulphonic acid in methanol. Finally, the protecting group is removed by hydrogenation using, for example, palladium on charcoal in the presence of ethanol under an atmosphere of hydrogen.

excessive bone or cartilage loss, including osteoporosis, gingival disease including gingivitis and periodontitis, arthritis, more specifically, osteoarthritis and rheumatoid arthritis, Paget's disease; hypercalcemia of malignancy, and metabolic bone disease.

- Metastatic neoplastic cells also typically express high levels of proteolytic enzymes that degrade the surrounding matrix, and certain tumors and metastatic neoplasias may be effectively treated with the compounds of this invention.

- The present invention also provides methods of treatment of diseases caused by pathological levels of proteases, particularly cysteine and serine proteases, more particularly cysteine proteases, even more particularly as inhibitors of cysteine proteases of the papain superfamily, yet more particularly cysteine proteases of the cathepsin family, which methods comprise administering to an animal, particularly a mammal, most particularly a human in need thereof a compound of the present invention. The present invention especially provides methods of treatment of diseases caused by pathological levels of cathepsin K, which methods comprise administering to an animal, particularly a mammal, most particularly a human in need thereof, an inhibitor of cathepsin K, including a compound of the present invention. The present invention particularly provides methods for treating diseases in which cysteine proteases are implicated, including infections by pneumocystis carinii, trypanoma cruzi, trypanoma brucei, and Crithidia fusciculata; as well as in schistosomiasis, malaria, tumor metastasis, metachromatic leukodystrophy, muscular dystrophy, amyotrophy, and especially diseases in which cathepsin K is implicated, most particularly diseases of excessive bone or cartilage loss, including osteoporosis, gingival disease including gingivitis and periodontitis, arthritis, more specifically, osteoarthritis and rheumatoid arthritis, Paget's disease, hypercalcemia of malignancy, and metabolic bone disease.

- This invention further provides a method for treating osteoporosis or inhibiting bone loss which comprises internal administration to a patient of an effective amount of a compound of formula (I), alone or in combination with other inhibitors of bone resorption, such as bisphosphonates (i.e., allendronate), hormone replacement therapy, anti-estrogens, or calcitonin. In addition, treatment with a compound of this invention and an anabolic agent, such as bone morphogenic protein, iproflavone, may be used to prevent bone loss or to increase bone mass.

- In accordance with this invention, an effective amount of the compounds of formula (I) is administered to inhibit the protease implicated with a particular condition or disease. Of course, this dosage amount will further be modified according to the type of administration of the compound. For example, "effective amount" for acute therapy, parenteral administration of a compound of formula (I) is preferred. An intravenous infusion of the compound in 5% dextrose in water or normal saline, or a similar

formulation with suitable excipients, is most effective, although an intramuscular bolus injection is also useful. Typically, the parenteral dose will be about 0.01 to about 100 mg/kg; preferably between 0.1 and 20 mg/kg, in a manner to maintain the concentration of drug in the plasma at a concentration effective to inhibit cathepsin K. The

5 compounds are administered one to four times daily at a level to achieve a total daily dose of about 0.4 to about 400 mg/kg/day. The precise amount of an inventive compound which is therapeutically effective, and the route by which such compound is best administered, is readily determined by one of ordinary skill in the art by comparing the blood level of the agent to the concentration required to have a therapeutic effect.

10 Prodrugs of compounds of the present invention may be prepared by any suitable method. For those compounds in which the prodrug moiety is a ketone functionality, specifically ketals and/or hemiacetals, the conversion may be effected in accordance with conventional methods.

The compounds of this invention may also be administered orally to the patient, 15 in a manner such that the concentration of drug is sufficient to inhibit bone resorption or to achieve any other therapeutic indication as disclosed herein. Typically, a pharmaceutical composition containing the compound is administered at an oral dose of between about 0.1 to about 50 mg/kg in a manner consistent with the condition of the patient. Preferably the oral dose would be about 0.5 to about 20 mg/kg.

20 No unacceptable toxicological effects are expected when compounds of the present invention are administered in accordance with the present invention.

The compounds of this invention may be tested in one of several biological assays to determine the concentration of a compound which is required to have a given pharmacological effect.

25

Determination of cathepsin K proteolytic catalytic activity

All assays for cathepsin K were carried out with human recombinant enzyme. Standard assay conditions for the determination of kinetic constants used a fluorogenic peptide substrate, typically Cbz-Phe-Arg-AMC, and were determined in 100 mM Na acetate at pH 5.5 containing 20 mM cysteine and 5 mM EDTA. Stock substrate solutions were prepared at concentrations of 10 or 20 mM in DMSO with 20 μ M final substrate concentration in the assays. All assays contained 10% DMSO. Independent experiments found that this level of DMSO had no effect on enzyme activity or kinetic constants. All assays were conducted at ambient temperature. Product fluorescence (excitation at 360 nM; emission at 460 nM) was monitored with a Perceptive Biosystems Cytofluor II fluorescent plate reader. Product progress curves were generated over 20 to 30 minutes following formation of AMC product.

Inhibition studies

Potential inhibitors were evaluated using the progress curve method. Assays were carried out in the presence of variable concentrations of test compound. Reactions were initiated by addition of enzyme to buffered solutions of inhibitor and substrate. Data analysis was conducted according to one of two procedures depending on the appearance of the progress curves in the presence of inhibitors. For those compounds whose progress curves were linear, apparent inhibition constants ($K_{i,app}$) were calculated according to equation 1 (Brandt *et al.*, *Biochemistry*, 1989, 28, 140):

$$v = V_m A / [K_d (1 + I/K_{i,app}) + A] \quad (1)$$

where v is the velocity of the reaction with maximal velocity V_m , A is the concentration of substrate with Michaelis constant of K_d , and I is the concentration of inhibitor.

For those compounds whose progress curves showed downward curvature characteristic of time-dependent inhibition, the data from individual sets was analyzed to give k_{obs} according to equation 2:

$$[AMC] = v_{ss} t + (v_0 - v_{ss}) [1 - \exp(-k_{obs}t)] / k_{obs} \quad (2)$$

where $[AMC]$ is the concentration of product formed over time t , v_0 is the initial reaction velocity, and v_{ss} is the final steady state rate. Values for k_{obs} were then analyzed as a linear function of inhibitor concentration to generate an apparent second order rate constant ($k_{obs} / \text{inhibitor concentration}$ or $k_{obs} / [I]$) describing the time-

dependent inhibition. A complete discussion of this kinetic treatment has been fully described (Morrison *et al.*, *Adv. Enzymol. Relat. Areas Mol. Biol.*, 1988, 61, 201).

- One skilled in the art would consider any compound with a K_i of less than 50 micromolar to be a potential lead compound. Preferably, the compounds used in the method of the present invention have a K_i value of less than 1 micromolar. Most preferably, said compounds have a K_i value of less than 100 nanomolar. 4-(*R,S*)-Amino-N-[(8-quinolinesulfonyl)-*S*-leucine]-3-tetrahydrofuran-3-one, a compound of formula (I), has a K_i value that is greater than 10 micromolar.

10 Human Osteoclast Resorption Assay

- Aliquots of osteoclastoma-derived cell suspensions were removed from liquid nitrogen storage, warmed rapidly at 37°C and washed x1 in RPMI-1640 medium by centrifugation (1000 rpm, 5 min at 4°C). The medium was aspirated and replaced with murine anti-HLA-DR antibody, diluted 1:3 in RPMI-1640 medium, and incubated for 30 minutes on ice. The cell suspension was mixed frequently.

The cells were washed x2 with cold RPMI-1640 by centrifugation (1000 rpm, 5 min at 4°C) and then transferred to a sterile 15 mL centrifuge tube. The number of mononuclear cells were enumerated in an improved Neubauer counting chamber.

- Sufficient magnetic beads (5 / mononuclear cell), coated with goat anti-mouse IgG, were removed from their stock bottle and placed into 5 mL of fresh medium (this washes away the toxic azide preservative). The medium was removed by immobilizing the beads on a magnet and is replaced with fresh medium.

- The beads were mixed with the cells and the suspension was incubated for 30 minutes on ice. The suspension was mixed frequently. The bead-coated cells were immobilized on a magnet and the remaining cells (osteoclast-rich fraction) were decanted into a sterile 50 mL centrifuge tube. Fresh medium was added to the bead-coated cells to dislodge any trapped osteoclasts. This wash process was repeated x10. The bead-coated cells were discarded.

- The osteoclasts were enumerated in a counting chamber, using a large-bore disposable plastic pasteur pipette to charge the chamber with the sample. The cells were pelleted by centrifugation and the density of osteoclasts adjusted to 1.5×10^4 /mL in EMEM medium, supplemented with 10% fetal calf serum and 1.7g/litre of sodium bicarbonate. 3 mL aliquots of the cell suspension (per treatment) were decanted into 15 mL centrifuge tubes. These cells were pelleted by centrifugation. To each tube 3 mL of the appropriate treatment was added (diluted to 50 μ M in the EMEM medium). Also included were appropriate vehicle controls, a positive control (87MEM1 diluted to 100

ug/mL) and an isotype control (IgG2a diluted to 100 ug/mL). The tubes were incubated at 37°C for 30 minutes.

0.5 mL aliquots of the cells were seeded onto sterile dentine slices in a 48-well plate and incubated at 37°C for 2 hours. Each treatment was screened in quadruplicate.

- 5 The slices were washed in six changes of warm PBS (10 mL / well in a 6-well plate) and then placed into fresh treatment or control and incubated at 37°C for 48 hours. The slices were then washed in phosphate buffered saline and fixed in 2% glutaraldehyde (in 0.2M sodium cacodylate) for 5 minutes, following which they were washed in water and incubated in buffer for 5 minutes at 37°C. The slices were then washed in cold water
10 and incubated in cold acetate buffer / fast red garnet for 5 minutes at 4°C. Excess buffer was aspirated, and the slices were air dried following a wash in water.

The TRAP positive osteoclasts were enumerated by bright-field microscopy and were then removed from the surface of the dentine by sonication. Pit volumes were determined using the Nikon/Lasertec ILM21W confocal microscope.

15

Examples

In the following synthetic examples, unless otherwise indicated, all of the starting materials were obtained from commercial sources. Without further elaboration, it is believed that one skilled in the art can, using the preceeding description, utilize the
20 present invention to its fullest extent. These Examples are given to illustrate the invention, not to limit its scope. Reference is made to the claims for what is reserved to the inventors hereunder.

Example 1

25

Preparation of 4-(R,S)-Amino-N-[(3,4-methylenedioxybenzoyl)-S-leucine]-tetrahydrofuran-3-one

a) *trans*-4-Azido-3-hydroxytetrahydrofuran

30

3,4-Epoxytetrahydrofuran (9 g, 105 mmol) was added to a stirred solution of sodium azide (27 g, 415 mmol) and ammonium chloride (9 g, 159 mmol) in aqueous methanol (95%, 200 mL). The reaction was heated to 75°C and stirred for 20 hours. The reaction was cooled, filtered and evaporated under reduced pressure. The residue was diluted with water and
35 extracted with ethyl acetate, dried and evaporated under reduced pressure to afford the title compound as a colourless oil, 10 g, 74% yield.

¹H NMR δ (CDCl₃) 4.32 (m, 1H), 4.09 (dd, 1H, J = 4.8, 9.9 Hz), 3.99 (dd, 1H, J = 4.3, 10.1 Hz), 3.94 (m, 1H), 3.81 (dd, 1H, J = 2.1, 9.9 Hz), 3.73 (dd, 1H, J = 1.8, 10.1 Hz), 2.72 (d, 1H, J = 4.6 Hz).

5 b) *trans*-4-Amino-3-hydroxytetrahydrofuran hydrochloride

A mixture of *trans*-4-azido-3-hydroxytetrahydrofuran (10 g, 77 mmol) and 10% palladium on charcoal (1 g) in ethanol (150 ml) was stirred under an atmosphere of hydrogen (35 psi) for 12 hours. The mixture was filtered and treated with 100 ml of ethanolic HCl to afford, after evaporation under reduced pressure, the title compound as a brown solid, 10.5 g, 97% yield. m.p. 132 °C

¹H NMR δ (d, DMSO) 8.37 (s, 3H), 4.13 (m, 1H), 3.84 (dd, 1H, J = 4.9 and 14.3), 3.76 (dd, 1H, J = 5.5, 10.0 Hz), 3.58 (dd, 1H, J = 2.7, 10.0 Hz), 3.34 (m, 3 H).

15 c) *trans*-4-(*R,S*)-Amino-N-[(*tert*-butoxycarbonyl)-*S*-leucine]-3-hydroxytetrahydrofuran

Trimethylacetyl chloride (3.5 ml, 29 mmol) was added to a stirred solution of N-Boc-L-leucine (7.3 g, 31 mmol) and diisopropylethylamine (9 ml, 52 mmol) in dichloromethane (200 ml). After 1 h, *trans*-4-amino-3-hydroxytetrahydrofuran. HCl (4 g, 28 mmol) was added and the mixture was allowed to stir overnight. The reaction mixture was poured into water and extracted with dichloromethane. The combined organic layers were washed with 0.5N HCl, saturated sodium hydrogen carbonate, brine and dried. Evaporation under reduced pressure afforded the title compound as a yellow foam, 5 g, 44% yield.

¹H NMR δ (CDCl₃) 8.08 (d, 0.5H, J = 4.8 Hz), 7.89 (d, 0.5H, J = 7.4 Hz), 6.20 (d, 0.5H, J = 8.3 Hz), 6.09 (δ, 0.5H, J = 8.7 Hz), 4.81 (d, 1H, J = 16.0Hz), 4.40 (m, 2H), 4.20 (m, 2H), 3.77 (m, 2H), 1.60 (m, 3H), 1.50 (s, 9H), 0.92 (m, 6H).

d) *trans*-4-(*R,S*)-Amino-N-(*S*-leucine)-3-hydroxytetrahydrofuran. TFA salt

trans-4-Amino-N-[(*tert*-butoxycarbonyl)-*S*-leucine]-3-hydroxytetrahydrofuran (2.5 g, 8.0 mmol) was added to a stirred solution of 20% trifluoroacetic acid in dichloromethane (100 ml). After 2 hours, the reaction mixture was evaporated under reduced pressure to afford the title compound as a white gum, 2.6 g, 100% yield.

¹H NMR δ (MeOD) 4.18 (m, 2H), 4.08 (m, 2H), 3.97 (m, 2H), 3.86 (apparent t, 2H, J = 7.1 Hz), 3.69 (dd, 2H, J = 1.6, 7.4 Hz), 1.68 (m, 3H), 0.99 (d, 6H, J = 2.1 Hz).

e) *trans*-4-(*R,S*)-Amino-N-[(3,4-methylenedioxybenzoyl)-*S*-leucine]-3-hydroxytetrahydrofuran

- 5 Piperonyl chloride (400 mg, 2.2 mmol) was added to a stirred solution of *trans*-4-amino-N-(*S*-leucine)-3-hydroxytetrahydrofuran.TFA salt (380 mg, 1.1 mmol) in saturated sodium hydrogen carbonate (10 ml) and 1,4 dioxane (10 ml). The reaction was allowed to stir for 1 hour then diluted with ether. The organic layer was washed with 1N hydrochloric acid, sodium hydrogen carbonate, brine and dried. Evaporation of the solvent gave the title compound as a white foam, 250 mg, 60% yield.
- 10 ¹H NMR δ (CDCl₃) 8.23 (d, 0.5H, J = 4.8 Hz), 8.15 (d, 0.5H, J = 7.4 Hz), 7.85 (d, 0.5H, J = 7.6 Hz), 7.3 (m, 2H), 6.66 (dd, 1H, J = 8.1, 11.5 Hz), 5.92 (d, 2H, J = 6.2 Hz), 4.78 (m, 1H), 4.50 (s, 1H), 4.2 (s, 1H), 4.08-3.75 (m, 4H), 3.74-3.48 (m, 3H), 1.82-1.48 (m, 3H), 0.90 (m, 6H). MS calcd for (C₁₈H₂₃N₂O₅ + H)⁺: 365. Found: 365.

- 15 f) 4-(*R,S*)-Amino-N-[(3,4-methylenedioxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one

- Dess-Martin periodinane (500 mg, 1.2 mmol) was added to a stirred solution of *trans*-4-(*R,S*)-amino-N-[(3,4-methylenedioxybenzoyl)-*S*-leucine]-3-hydroxytetrahydrofuran (220 mg, 0.60 mmol) in dichloromethane (10 ml). After 1 hour, ether was added followed by sodium thiosulfate (570 mg, 3.6 mmol). After an additional 15 minutes, the reaction was washed with saturated sodium hydrogen carbonate, brine and dried. Evaporation of the solvent gave the title compound as a white foam, 200 mg, 100% yield.
- 20 ¹H NMR δ (CDCl₃) 8.14 (d, 0.5H, J = 6.2 Hz), 7.90 (d, 0.5H, J = 5.9 Hz), 7.54 (d, 0.5H, J = 7.3 Hz), 7.46 (d, 0.5H, J = 5.1 Hz), 7.23 (d, 1H, J = 6.6 Hz), 7.14 (s, 1H), 6.67 (m, 1H), 5.93 (s, 2H), 4.73 (m, 1H), 4.37-3.71 (m, 5H), 1.68 (m, 3H), 0.85 (m, 6H).
- 25 MS calcd for (C₁₈H₂₂N₂O₆ + H)⁺: 363. Found: 363.

Example 2

- 30 Preparation of 4-(*R,S*)-Amino-N-[(3,4-dichlorobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one

- Following the procedure of Example 1(e-f) except substituting 3,4-dichlorobenzoyl chloride for piperonyl chloride the title compound was prepared: ¹H NMR δ (MeOD) 8.01 (d, 1H, J = 2.0 Hz), 7.75 (dd, 1H, J = 2.0, 8.3 Hz), 7.61 (d, 1H, J = 8.3 Hz), 4.60 (m, 1H), 4.50-3.84 (m, 5H), 1.66 (m, 3H), 1.00 (m, 6H).
- 35 MS calcd for (C₁₇Cl₂H₂₀N₂O₄ + H)⁺: 387 & 389, Found: 387 & 389.

Example 3Preparation of 4-(*R,S*)-Amino-N-[(2-quinolinecarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

- 5 Following the procedure of Example 1(e-f) except substituting 2-quinolinecarbonyl chloride for piperonyl chloride, the title compound was prepared as a white foam: ^1H NMR δ (CDCl_3) 8.57 (dd, 1H, $J = 1.5, 5.1$ Hz), 8.20 (m, 3H), 7.77 (m, 2H), 7.62 (m, 1H), 7.40 (d, 1H, $J = 6.2$ Hz), 4.80 (m, 1H), 4.57 (m, 1H), 4.30 (m, 1H), 4.26-3.87 (m, 3H), 1.96-1.71 (m, 3H), 1.00 (m, 6H).
- 10 MS calcd for ($\text{C}_{30}\text{H}_{37}\text{N}_4\text{O}_4 + \text{H}$) $^+$: 370. Found: 370.

Example 4Preparation of 4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

- 15 Following the procedure of Example 1 except substituting N-CBZ-leucine for N-BOC-leucine the title compound was prepared as a white foam: ^1H NMR δ (CDCl_3) 7.31 (m, 5H), 5.71 (app dd, 1H, $J = 8.4, 14.8$ Hz), 5.06 (m, 2H), 4.50 (dd, 1H, $J = 8.9, 18.0$ Hz), 4.30-3.88 (m, 4H), 3.80 (app t, 1H, $J = 9.6$ Hz), 1.64 (m, 3H), 0.91 (m, 6H).
- 20 MS calcd for ($\text{C}_{18}\text{H}_{23}\text{N}_2\text{O}_5 + \text{H}$) $^+$: 349. Found: 349.

Example 5Preparation of 4-(*R,S*)-Amino-N-[(3,4-methylenedioxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one

- 25 Following the procedure of Example 1 except substituting 3,4-methylenedioxybenzoyl chloride for piperonyl chloride, the title compound was prepared.
- 30

Example 6Preparation of 4-(*R,S*)-Amino-N-[(8-quinolinecarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

5

Following the procedure of Example 1(e-f) except substituting 8-quinolinecarbonyl chloride for piperonyloyl chloride, the title compound was prepared as a white foam: m.p. 123 °C (HCl salt).

¹H NMR δ (CDCl₃) 11.54 (m, 1H), 8.84 (d, 1H, J = 1.5 Hz), 8.65 (d, 1H, J = 7.2 Hz), 8.17 (d, 1H, J = 8.0 Hz), 7.86 (d, 1H, J = 8.0 Hz), 7.7 (s, 1H), 7.54 (t, 1H, J = 7.7 Hz), 7.40 (dd, 1H, J = 3.8, 7.7 Hz), 4.78 (dd, 1H, J = 7.5, 13.6 Hz), 4.47 (dd, 1H, J = 8.6, 13.6 Hz), 4.18 (m, 1H), 4.16-3.79 (m, 3H), 1.88 (m, 3H), 0.88 (m, 6H).
MS calcd for (C₂₀H₂₃N₃O₄ + H)⁺: 370. Found: 370.

15

Example 7Preparation of 4-(*R,S*)-Amino-N-[(8-quinolinesulfonyl)-*S*-leucine]-tetrahydrofuran-3-one

20

Following the procedure of Example 1(e-f) except substituting 8-quinolinesulfonyl chloride for piperonyloyl chloride, the title compound was prepared as a brown solid: m.p. 98 °C (HCl salt).

¹H NMR δ (CDCl₃) 9.07 (dd, 1H, J = 1.7 and 4.3 Hz), 8.40 (m, 1H), 8.30 (m, 1H), 8.00 (m, 1H), 7.66 (m, 2H), 7.11 (d, 0.5H, J = 5.9 Hz), 6.96 (d, 0.5H, J = 5.7 Hz), 4.51 (t, 0.5H, J = 8.8 Hz), 4.39 (t, 0.5H, J = 8.7), 4.12-3.71 (m, 4.5H), 3.53 (dd, 0.5H, J = 1.0, 5.9 Hz), 1.43 (m, 2H), 0.64 (m, 3H), 0.33 (m, 3H).
MS calcd for (C₁₉H₂₃N₃O₅S + H)⁺: 406. Found: 406.

30

Example 8Preparation of 4-(*R,S*)-Amino-N-[(4-methyl-3-pyridinyl)carbonyl]-*S*-leucine]-tetrahydrofuran-3-one

Following the procedure of Example 1(e-f) except substituting (4-methyl-3-pyridinyl)carbonyl chloride for piperonyloyl chloride, the title compound was prepared.

35

Example 9Preparation of 4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-2,2-dibenzyl-tetrahydrofuran-3-one

5

Sodium methoxide (140 mg, 2.6 mmol) was added to a stirred solution of 4-(*R,S*)-amino-N-[(benzyloxycarbonyl)-*S*-leucine]-tetrahydrofuran-3-one (300 mg, 0.9 mmol) and benzyl bromide (0.4 ml, 3.4 mmol) in methanol (5 ml). After 12 hours, the reaction was poured into ether (100 ml) and washed with water, brine and dried.

- 10 Evaporation under reduced pressure and purification by flash column chromatography (30% ethyl acetate-hexane) afforded the title compound as a colorless gum, 250 mg, 55% yield.

¹H NMR δ (CDCl₃) 7.34-7.25 (m, 10 H), 7.24-7.0 (brs, 5 H), 6.56 (s, 0.5 H), 6.37 (s, 0.5 H), 5.18 (d, 1 H, J = 11.5 Hz), 5.07 (m, 2 H), 4.48-4.42 (m, 1 H), 4.20-3.98 (m, 4 H), 3.12 (dd, 1 H, J = 12.5 and 12.5 Hz), 2.88 (dd, 1 H, J = 12.5 and 12.5 Hz), 1.72-1.32 (m, 3 H), 0.92-0.76 (m, 6 H).

- 15

MS calcd for (C₃₂H₃₆N₂O₅ = C₇H₇)⁺: 439. Found: 439

Example 10

20

Preparation of 4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

Following the procedure of Example 1(e-f) except substituting

- 25 benzo[b]thiophen-2-ylcarbonyl chloride for piperonyl chloride, the title compound was prepared.

¹H NMR δ (CDCl₃) 8.33 (d, 0.5H, J = 6.6Hz), 8.00 (m, 1H), 7.78 (m, 4H), 7.38 (m, 2.5H), 4.87 (m, 1H), 4.63-3.88 (m, 5H), 1.88 (m, 3H), 1.00 (m, 6H).

MS calcd for (C₁₈H₂₀N₂O₄S - H)⁺: 373, Found: 373.

30

Example 11Preparation of 4-(*R,S*)-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one

35

Following the procedure of Example 1(e-f) except substituting 3,4-dimethoxybenzoyl chloride for piperonyl chloride, the title compound was prepared.

¹H NMR δ (CDCl₃) 7.58 (d, 0.5H, J = 6.7Hz), 7.47 (d, 0.5H, J = 6.2Hz), 7.33 (m, 2H), 7.04 (d, 0.5H, J = 8.0Hz), 6.92 (d, 0.5H, J = 8.0Hz), 6.83 (d, 0.5H, J = 8.4Hz), 4.77 (m, 1H), 4.53-3.67 (m, 5H), 1.68 (m, 3H), 0.85 (m, 6H).

MS calcd for (C₁₉H₂₆N₂O₄ - H)⁺: 377, Found: 377.

5

Example 12

Preparation of 4-(R,S)-Amino-N-[(indole-6-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-one

10

Following the procedure of Example 1(e-f) except substituting indole-6-ylcarbonyl chloride for piperonyl chloride, the title compound was prepared.

¹H NMR δ (d₆ DMSO) 8.57-7.48 (m, 7H), 6.43 (s, 1H), 4.48-3.55 (m, 6H), 1.80-1.48 (m, 3H), 0.98-0.82 (m, 6H).

15 MS calcd for (C₁₉H₂₃N₂O₄ - H)⁺: 356, Found: 356.

Example 13

Preparation of 4-(R,S)-Amino-N-[(benzofuran-2-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-one

20

Following the procedure of Example 1(e-f) except substituting benzofuran-2-ylcarbonyl chloride for piperonyl chloride, the title compound was prepared.

¹H NMR δ (CDCl₃) 8.00-7.25 (m, 7H), 4.90-4.78 (m, 1H), 4.53-4.48 (m, 1H), 4.38-4.21 (m, 1H), 4.25-3.92 (m, 3H), 1.88 (m, 3H), 1.68 (m, 3H), 0.97 (m, 6H).

25

MS calcd for (C₁₉H₂₂N₂O₅ - H)⁺: 357, Found: 357.

Example 14

Preparation of 4-(R,S)-Amino-N-[(5-aminobenz[b]thiophen-2-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-one

30

Following the procedure of Example 1(e-f) except substituting 5-aminobenz[b]thiophen-2-ylcarbonyl chloride for piperonyl chloride, the title compound was prepared.

35

¹H NMR δ (CDCl₃) 8.00-7.71 (m, 1H), 7.60 (d, 1H, J = 3.3Hz), 7.53 (dd, 1H, J = 8.5 and 4.0 Hz), 7.47-6.74 (m, 3H), 4.77 (m, 1H), 4.43 (m, 1H), 4.38-3.50 (m, 4H) 1.77 (m, 3H), 0.85 (m, 6H).

MS calcd for (C₁₆H₂₃N₃O₂S + H)⁺: 390, Found: 390.

5

Example 15

Preparation of Cyclic Alkoxyketones by Solid Supported Synthesis

10 a) 3-*trans*-Hydroxy-4-benzyloxycarbonylamino-tetrahydrofuran

Benzylchloroformate (20ml) was added dropwise to a stirred solution of *trans*-4-amino-3-hydroxytetrahydrofuran (5g, 20.6mmol) in dioxane (100ml) containing 10% aqueous sodium carbonate (200ml). After 3 hours, the mixture was concentrated to remove the dioxane and then extracted with EtOAc. The combined organic layers were washed with saturated sodium bicarbonate, brine and dried (MgSO₄). Evaporation under reduced pressure and purification of the residue by column chromatography afforded the title compound as white crystals, 8.00g, 70%.

¹H NMR δ (CDCl₃) 7.35 (s, 5H), 5.30 (s, 2H), 4.91 (br s, 1H), 4.32 (br s, 1H), 4.12-4.00 (m, 3H), 3.71-3.62 (m, 2H), 2.72 (s, 1H).

20 b) 4-Benzyloxycarbonylamino-tetrahydrofuran-3-one

A solution of bleach (100ml), containing sodium bicarbonate (7.34g), was added dropwise to a rapidly stirred mixture of 3-hydroxy-4-benzyloxycarbonylamino-tetrahydrofuran (21g, 88 mmol), sodium bromide (9.4g), TEMPO (50mg) in EtOAc (140ml), toluene (140ml) and water (40ml). After a persistent orange colour developed the mixture was extracted with EtOAc and the combined organic layers were washed with saturated sodium bicarbonate, brine and dried (MgSO₄). Evaporation under reduced pressure and purification of the residue by column chromatography afforded the title compound as white crystals, 18g, 87%.

¹H NMR δ (CDCl₃) 7.35 (s, 5H), 5.30 (s, 1H), 5.11 (s, 2H), 4.70 (app t, 1H, J = 8.9Hz), 4.36-4.17 (m, 2H), 3.96-3.76 (2H).

35 c) 3,3-Dimethoxy-4-benzyloxycarbonylamino-tetrahydrofuran

Trimethylorthoformate (29ml) was added dropwise to a refluxing solution of 4-benzyloxycarbonylamino-tetrahydrofuran-3-one (18g, 78mmol) and PTSA(500mg) in MeOH (100ml). After 3 hours, the reaction mixture was filtered and concentrated to afford, after column chromatography, the title compound as a yellow oil, 16.2g, 76%.

40

¹H NMR δ (CDCl₃) 7.35 (s, 5H), 5.30 (s, 1H), 5.11 (s, 2H), 4.70 (app t, 1H, J = 8.9Hz), 4.36-4.17 (m, 2H), 3.96-3.76 (2H).

d) 3,3-Dimethoxy-4-amino-tetrahydrofuran

5

A mixture of 3,3-Dimethoxy-4-benzyloxycarbonylamino-tetrahydrofuran (16g, 57mmol) and 10% palladium on charcoal (2g) in ethanol (200ml) was stirred under an atmosphere of hydrogen (50psi) for 12 h. The mixture was filtered and concentrated to afford the title compound as a yellow oil, 8g, 100%.

10 ¹H NMR δ (CDCl₃) 7.04 (s, 5H), 4.22-4.04 (m, 5H), 3.83-3.69 (m, 2H), 3.36 (s, 3H), 3.33 (s, 3H).

e) SPS using Ellman Linker

15 *Step A:*

Sodium triacetoxymethylborohydride (10 equiv) was added to a stirred solution of Ellmans resin (ref: C.G. Boojamra, K.M. Burow, L.A. Thompson and J.A. Ellman, *J. Org. Chem.*, 1997, 62, 1240) in DMF containing 1% HOAc. After 5 minutes, the α -amino acid, methyl ester (10 equiv) was added and the mixture was shaken for 1 hour.

20 The resin was then washed with DMF (x7), CH₂Cl₂ (x7), ether (x2) and dried to a constant weight.

Step B:

NMP was added to a mixture of the above resin, carboxylic acid (10 equiv) and EDC (10 equiv). The mixture was then shaken for 3 hours then washed with DMF (x3), CH₂Cl₂ (x3), MeOH (x2) and ether (x2). The resin was then resubjected to the above reaction conditions and again washed after 3 hours.

25

Step C:

Potassium trimethylsilanoate (10 equiv) was added to a shaken mixture of the above resin in THF. After 18 hours, the resin was washed with 5% citric acid in THF (x2), THF (x2), THF-H₂O (x2), H₂O (x2), THF-H₂O (x2) and finally THF (x2).

30

Step D:

3,3-Dimethoxy-4-amino-tetrahydrofuran (3 equiv) was added to a mixture of the above resin and EDC (3 equiv) in NMP. After 3 hours, the resin was washed with DMF (x7), CH₂Cl₂ (x7) and ether (x2). The resin was then resubjected to the reaction conditions for a further 3 h, then again washed as above.

35

40

Step E: Cleavage.

A mixture of 7:2:1 TFA/CH₂Cl₂/H₂O was added to the above resin. After 2 hours, the mixture was filtered and the resin was further washed with CH₂Cl₂. Removal of the solvent afforded the desired tetrahydrofuran-3-one.

Example 16Preparation of 4-(R,S)-Amino-N-[(5-chlorobenzofuran-2-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-one

Following the general details of Example 15, using the appropriate amino acid methyl ester and carboxylic acid reagents consistent with the final product, the title compound was prepared.

¹H NMR δ (CDCl₃) 7.62-7.05 (m, 6H), 4.81-4.64 (m, 1H), 4.62-4.54 (m, 1H), 4.43-3.81 (m, 4H), 1.90-1.60 (m, 3H), 1.08-0.81 (m, 6H)

MS calcd for (C₁₉H₂₁N₂O₅Cl + H)⁺: 393, Found: 393.

Example 17Preparation of 4-(R,S)-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-one

Following the general details of Example 15, using the appropriate amino acid methyl ester and carboxylic acid reagents consistent with the final product, the title compound was prepared.

¹H NMR δ (CDCl₃) 7.65-6.84 (m, 6H), 4.83-4.54 (m, 1H), 4.53-4.48 (m, 1H), 4.46-3.68 (m, 4H), 1.90-1.62 (m, 3H), 1.08-0.81 (m, 6H).

MS calcd for (C₂₀H₂₃N₂O₆ + H)⁺: 389, Found: 389.

Example 18Preparation of 4-(R,S)-Amino-N-[(4-bromobenzoyl)-S-leucine]-tetrahydrofuran-3-one

Following the general details of Example 15, using the appropriate amino acid methyl ester and carboxylic acid reagents consistent with the final product, the title compound was prepared.

¹H NMR δ (CDCl₃) 7.64 (App d, 2H, J = 8.5Hz), 7.60 (App d, 2H, J = 8.5Hz), 4.81-4.67 (m, 1H), 4.62-4.48 (m, 1H), 3.36-4.19 (m, 1H), 4.18-3.78 (m, 3H), 1.81-1.59 (m, 3H), 1.05-0.80 (m, 6H).

MS calcd for (C₁₇H₂₁N₂O₄Br)⁺: 397, Found: 397.

5

Example 19

Preparation of 4-(R,S)-Amino-N-[(4-bromobenzoyl)-S-leucine]-tetrahydrofuran-3-one

10 Following the general details of Example 15, using the appropriate amino acid methyl ester and carboxylic acid reagents consistent with the final product, the title compound was prepared.

¹H NMR δ (CDCl₃) 7.91 (m, 1H), 7.76-7.59 (m, 2H), 7.39-7.18 (m, 1H), 7.04-6.89 (m, 1H), 7.84-6.68 (m, 1H), 4.89-4.468 (m, 1H), 4.66-4.56 (m, 1H), 4.27-3.76 (m, 4H), 1.88-1.68 (m, 3H), 1.03-0.78 (m, 6H)

15

MS calcd for (C₁₇H₂₁N₂O₄Br)⁺: 397, Found: 397.

Example 20

20 Preparation of 4-(R,S)-Amino-N-[(5-chlorobenzoyl)-S-leucine]-tetrahydrofuran-3-one

Following the general details of Example 15, using the appropriate amino acid methyl ester and carboxylic acid reagents consistent with the final product, the title compound was prepared.

25

¹H NMR δ (CDCl₃) 7.88-7.67 (m, 5H), 7.48-7.38 (m, 1H), 4.81 (br d, 1H, J = 6.7Hz), 4.60 (app t, 1H, J = 8.8Hz), 4.43-4.30 (m, 1), 4.28-3.84 (m, 3H), 1.86-1.62 (m, 3H), 1.05-0.82 (m, 6H).

MS calcd for (C₁₉H₂₁N₂O₄SCl - H)⁺: 408, Found: 408.

30

Example 21

Preparation of 4-(R,S)-Amino-N-[(4-fluorobenzoyl)-S-leucine]-tetrahydrofuran-3-one

35 Following the general details of Example 15, using the appropriate amino acid methyl ester and carboxylic acid reagents consistent with the final product, the title compound was prepared.

^1H NMR δ (CDCl_3) 7.96 (s, 1H), 7.68-7.58 (m, 2H), 7.48-7.32 (m, 1H), 7.06 (dd, 1H, $J = 9.0\text{ Hz}$ and 9.0 Hz), 4.89-4.72 (m, 1H), 4.65-4.55 (app t, 1H, $J = 8.8\text{ Hz}$), 4.35 (app q, 1H, $J = 8.2\text{ Hz}$), 4.28-3.82 (m, 3H), 1.88-1.62 (m, 3H), 1.05-0.82 (m, 6H).
 MS calcd for $(\text{C}_{10}\text{H}_{11}\text{N}_3\text{O}_3\text{SF} + \text{H})^+$: 393, Found: 393.

5


Examples 22-66

Preparation of Cyclic Alkoxyketones




- 10 By analogous procedures to those described in either Example 1 or Example 15, using the appropriate amino acid and acid or acid chloride reagents consistent with the final products, the compounds of Table 1 were prepared. ^1H NMR spectra and/or mass spectra were consistent with the structures in Table 1.

15

Table 1



Example	R^3	R^1	synthesis method
22		3,4-difluorophenyl	soln
23	"	4-benzylpiperidin-1-yl	soln
24	"	4-benzylpiperazin-1-yl	soln
25	"	4-(3,4-methylenedioxybenzyl)piperazin-1-yl	soln
26	"	4-(tert-butoxycarbonyl)piperazin-1-yl	soln
27	"	piperazin-1-yl	soln
28	"	benzimidazol-5-yl	soln
29	"	6-quinolyl	soln
30	"	5-indolyl	soln
31	"	2-naphthyl	soln
32	"	2-pyridyl	soln
33	"	4-benzoyloxyphenyl	soln
34	"	3-benzoyloxyphenyl	soln
35	"	4-hydroxyphenyl	soln
36	"	5-nitrobenzo[b]thiophen-2-yl	soln
37	"	2-(thien-2-yl)ethen-1-yl	soln
38	"	4-methoxyphenyl	SPS

39	"	3-methoxyphenyl	SPS
40	"	7-ethoxybenzofuran-1-yl	SPS
41	"	5-nitrobenzofuran-1-yl	SPS
42	"	4-(2-methoxyphenyl)phenyl	SPS
43	"	3-(2-methoxyphenyl)phenyl	SPS
44	"	4-cyanophenyl	SPS
45	"	3-nitrophenyl	SPS
46	"	3-(dimethylaminoethyl)-4-methoxyphenyl	SPS
47	"	2-(2-chlorophenyl)ethen-1-yl	SPS
48	"	4-trifluoromethoxyphenyl	SPS
49	"	4-methanesulphonylphenyl	SPS
50	"	4-iodophenyl	SPS
51	"	4-chlorobenzo[b]thiophen-2-yl	SPS
52	"	5,6-dimethoxybenzo[b]thiophen-2-yl	SPS
53	"	5,6-methylenedioxybenzo[b]thiophen-2-yl	SPS
54	"	7-chlorobenzo[b]thiophen-2-yl	SPS
55		benzo[b]thiophen-2-yl	SPS
56	"	2-thienyl	SPS
57	"	3,4-dimethoxyphenyl	SPS
58	"	4-bromophenyl	SPS
59	"	quinolin-2-yl	SPS
60		2-thienyl	SPS
61	"	CH ₂	SPS
62		benzo[b]thiophen-2-yl	SPS
63	"	2-thienyl	SPS
64	"	3,4-dimethoxyphenyl	SPS
65	"	4-bromophenyl	SPS
66	"	quinolin-2-yl	SPS

Example 67Preparation of 4-(R,S)-Amino-N-[(benzyloxycarbonyl)-S-leucine]-tetrahydropyran-3-one

a) *trans*-4-amino-3-hydroxytetrahydropyran hydrochloride

Following the procedures of Examples 1(a) and 1(b), substituting 3,4-epoxytetrahydropyran for 3,4-epoxytetrahydrofuran, the title compound was prepared.

b) *trans*-4-(R,S)-amino-N-[(benzyloxycarbonyl)-S-leucine]-3-hydroxytetrahydropyran

Pivaloyl chloride (5.28 ml, 43 mmol) was added to a solution of N-carbobenzyloxy-L-leucine (12.48 g, 47 mmol) in dichloromethane (300 ml). After 1 hour, a mixture of *trans*-4-amino-3-hydroxytetrahydropyran hydrochloride (6g, 39mmol) and triethylamine (10.8 ml, 79 mmol) in dichloromethane (100 ml) was added and the mixture allowed to stir overnight. The reaction mixture was washed with 1N HCl, saturated sodium hydrogen carbonate and dried. Evaporation under reduced pressure afforded a pale oil. Purification by chromatography (ethyl acetate/hexane eluant) gave the *trans*-4-(R,S)-amino-N-[(benzyloxycarbonyl)-S-leucine]-3-hydroxytetrahydropyran as a white paste, 5.5 g, 39% yield:
 ^1H NMR δ (CDCl₃) 7.35 (s, 5H), 5.0 (m, 3H), 4.16-3.89 (m, 4H), 3.44-3.36 (m, 2H), 3.13 (t, 1H), 1.84-1.53 (m, 2H), 1.28-1.22 (m, 3H), 0.93 (m, 6H).

c) 4-(R,S)-amino-N-[(benzyloxycarbonyl)-S-leucine]tetrahydropyran-3-one

Following the procedure of Example 1(f), except substituting 4-(R,S)-amino-N-[(benzyloxycarbonyl)-S-leucine]-3-hydroxytetrahydropyran for 4-(R,S)-amino-N-[(3,4-methylenedioxybenzoyl)-S-leucine]-3-hydroxytetrahydrofuran, the title compound was prepared:

^1H NMR δ (d₆ DMSO) 8.15 (d, 1H), 7.39-7.29 (m, 5H), 5.02 (d, 2H), 4.63 (m, 1H), 4.14-4.10 (m, 2H), 3.97-3.83 (m, 3H), 2.10 (m, 1H), 1.92 (m, 1H), 1.61 (m, 1H), 1.45 (m, 1H), 0.95 (m, 6H).

MS calcd for (C₁₉H₂₆N₂O₅+H)⁺: 363. Found 363.

Example 68Preparation of 4-(R,S)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-S-leucine]-tetrahydropyran-3-one

5

a) *trans*-4-(R,S)-Amino-N-(S-leucine)-3-hydroxytetrahydropyran hydrochloride

A mixture of 4-(R,S)-amino-N-[(benzyloxycarbonyl)-S-leucine]-3-hydroxytetrahydropyran (2.8 g, 7.75 mmol) and 10% palladium on charcoal (300 mg) in ethanol (100 ml) was stirred under an atmosphere of hydrogen (50 psi) for 12 hours. The mixture was filtered and treated with ethereal HCl to afford, after evaporation under reduced pressure, the title compound as a brown solid, 1.40 g, 68% yield:

¹H NMR δ (CDCl₃) 8.30 (m, 2H), 8.02 (m, 1H), 4.06-3.92 (m, 4H), 3.50-3.35 (m, 2H), 3.12 (t, 1H), 1.89-1.54 (m, 2H), 1.23 (m, 3H), 0.93 (m, 6H).

15

b) *trans*-4-(R,S)-amino-N-[(benzothiophene-2-carbonyl)-S-leucine]-3-hydroxytetrahydropyran

Benzothiophene-2-carbonyl chloride (442 mg, 2.25 mmol) was added to a solution of *trans*-4-(R,S)-amino-N-(S-leucine)-3-hydroxytetrahydropyran (133 mg, 0.5 mmol) in dioxan (7ml) and saturated sodium hydrogen carbonate (7 ml). After 30 min, the reaction mixture was diluted with ethyl acetate, the organic layer washed with saturated sodium hydrogen carbonate, dried and evaporated under reduced pressure to give a white solid. Purification by chromatography (ethyl acetate/hexane eluant) gave the *trans*-4-(R,S)-amino-N-[(benzothiophene-2-carbonyl)-S-leucine]-3-hydroxytetrahydropyran as a white solid, 160 mg, 84%.

25

c) 4-(R,S)-Amino-N-[(benzothiophene-2-carbonyl)-S-leucine]-tetrahydropyran-3-one

Following the procedure of Example 1(f) except substituting *trans*-4-(R,S)-amino-N-[(benzothiophene-2-carbonyl)-S-leucine]-3-hydroxytetrahydropyran for 4-(R,S)-amino-N-[(3,4-methylenedioxybenzoyl)-S-leucine]-3-hydroxytetrahydrofuran, the title compound was prepared:

¹H NMR δ (CDCl₃) 7.84-7.76 (m, 2H), 7.41 (m, 2H), 7.05 (m, 1H), 4.83-4.61 (m, 2H), 4.18-3.77 (m, 4H), 2.73-2.53 (m, 1H), 1.98-1.75 (m, 2H), 1.26 (m, 3H), 0.92 (m, 6H).

MS calcd for (C₂₀H₂₈N₂O₄S+H)⁺: 389. Found 389.

35

Example 69Preparation of 4-(*R,S*)-Amino-N-[(4-phenoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one

- 5 By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (CDCl₃, 250 MHz) 7.74 (d, 2 H, J = 10.9 Hz), 7.37 (dd, 2 H, J = 7.7 and 7.7 Hz), 7.20 (dd, 1 H, J = 7.5 and 7.5 Hz), 7.03 (d, 2 H, J = 7.7 Hz), 7.0 (d, 2 H, J = 7.7 Hz), 6.98-6.82 (m, 2 H), 4.83-4.68 (m, 1 H), 4.66-4.45 (m, 1 H), 4.34-3.70 (m, 4 H),

- 10 1.98-1.54 (m, 3 H), 1.08-0.78 (m, 6 H).

MS calcd for (C₂₃H₂₆N₂O₅ - H)⁺: 409. Found: 409.

Example 70

- 15 Preparation of 4-(*R,S*)-Amino-N-[(4-phenylbenzoyl)-*S*-leucine]-tetrahydrofuran-3-one

By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (CDCl₃, 250 MHz) 7.83 (d, 2 H, J = 8.2 Hz), 7.70-50 (m, 4 H), 7.48-32 (m, 4 H), 7.30-7.12 (m, 1 H), 6.88-6.72 (m, 1 H), 4.88-4.70 (m, 1 H), 4.65-52 (m, 1 H), 4.38-3.78 (m, 4 H), 1.92-1.60 (m, 3 H), 1.08-0.78 (m, 6 H).

MS calcd for (C₂₃H₂₆N₂O₄ - H)⁺: 393. Found: 393.

Example 71

25

Preparation of 4-(*R,S*)-Amino-N-[(6-trifluoromethylbenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

- By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (DMSO-d₆, 250 MHz) 8.82 (d, 1 H, J = 6.0 Hz), 8.43 (s, 1 H), 8.48-8.44 (m, 0.5 H), 8.34 (d, 0.5 H, J = 7.8 Hz), 8.23 (s, 1 H), 8.04 (d, 1 H, J = 8.4 Hz), 7.60 (dd, 1 H, J = 1.5 and 8.5 Hz), 4.49-4.38 (m, 1 H), 4.26-3.68 (m, 5 H), 1.70-1.36 (m, 3 H), 0.80 (d, 3 H, J = 6 Hz), 0.76 (d, 3 H, J = 6.0 Hz)

- 35 MS calcd for (C₂₉H₂₁F₃N₂O₅ - H)⁺: 443. Found: 443.

Example 72Preparation of 4-(*R,S*)-Amino-N-[(4-ethylbenzoyl)-*S*-leucine]-tetrahydrofuran-3-one

- 5 By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (CDCl₃, 250 MHz) 7.76 (d, 2 H, J = 8.0 Hz), 7.62-7.42 (m, 1 H), 7.22 (d, 2 H, J = 7.8 Hz), 7.04-6.80 (m, 1 H), 4.91-4.73 (m, 1 H), 4.61-4.45 (m, 1 H), 4.36-3.72 (m, 4 H), 2.68 (q, 2 H, J = 7.6 Hz), 1.88-1.58 (m, 3 H), 1.23 (t, 3 H, J = 7.6 Hz), 0.98-0.88 (m, 6 H)

MS calcd for (C₁₉H₂₉N₂O₄ - H)⁺: 345. Found: 345.

Example 73

- 15 Preparation of 4-(*R,S*)-Amino-N-[(4-(tert-butyl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one

By analogous methods to those detailed in Example 15, the title compound was prepared.

- 20 ¹H NMR δ (CDCl₃, 250 MHz) 7.76 (brd, 2 H, J = 7.5 Hz), 7.46 (brd, 2 H, J = 7.5 Hz), 6.92-6.76 (m, 2H), 4.88-4.68 (m, 1 H), 4.58-4.43 (m, 1 H), 4.37-3.71 (m, 4 H), 1.82-1.57 (m, 3 H), 1.32 (s, 9 H), 1.00-0.82 (m, 6 H).

MS calcd for (C₂₁H₃₃N₂O₄ + H)⁺: 375. Found: 375.

- 25 Example 74

Preparation of 4-(*R,S*)-Amino-N-[(5-methoxybenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

- 30 By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (CDCl₃, 250 MHz) 7.84-7.54 (m, 2 H), 7.20 (d, 1 H, J = 2.0 Hz), 7.10 (d, 1 H, J = 2.3 Hz), 7.06 (d, 1 H, J = 2.3 Hz), 6.78 (d, 1 H, J = 8.1 Hz), 4.82-4.68 (m, 1 H), 4.59 (appt, 1 H, J = 8.8 Hz), 4.49-3.61 (m, 4 H), 3.84 (s, 3 H), 1.82-1.58 (m, 3 H), 1.08-0.72 (m, 6 H).

MS calcd for (C₂₀H₂₈N₂O₅S - H)⁺: 403. Found: 403.

Example 75Preparation of 4-(*R,S*)-Amino-N-[(4-nitrobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

5

By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (CDCl₃, 250 MHz) 8.31 (s, 1 H), 7.96 (d, 1 H, J = 7.2 Hz), 7.62-7.18 (m, 4 H), 4.84 (appd, 1 H, J = 7.5 Hz), 4.70-3.72 (m, 5 H), 1.94-1.60 (m, 3 H), 1.11-0.80 (m, 6 H).

10

MS calcd for (C₁₉H₂₁N₃O₆S-NO₂)⁺: 373. Found: 373.

Example 7615 Preparation of 4-(*R,S*)-Amino-N-[(6-bromobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

By analogous methods to those detailed in Example 15, the title compound was prepared.

20 ¹H NMR δ (CDCl₃, 250 MHz) 8.03 (s, 1 H), 7.81-7.58 (m, 2 H), 7.54-7.40 (m, 1 H), 7.22-6.98 (m, 2 H), 4.74 (appd, 1 H, J = 7.5 Hz), 4.59 (appq, 1 H, J = 7.5 and 14.0 Hz), 4.48-3.74 (m, 4 H), 1.85-1.58 (m, 3 H), 1.10-0.78 (m, 6 H).

MS calcd for (C₁₉H₂₁BrN₃O₆S-H)⁺: 452. Found: 452.

25

Example 77Preparation of 4-(*R,S*)-Amino-N-[(5-bromobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

30 By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (CDCl₃, 250 MHz) 8.04-7.90 (m, 1 H), 7.88-7.18 (m, 5 H), 4.88-4.68 (m, 1 H), 4.68-4.52 (m, 1 H), 4.46-3.88 (m, 4 H), 1.92-1.52 (m, 3 H), 1.08-0.80 (m, 6 H).

MS calcd for (C₁₉H₂₁BrN₃O₆S-H)⁺: 452. Found: 452.

35

Example 78Preparation of 4-(*R,S*)-Amino-N-[(6-methoxybenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one

5

By analogous methods to those detailed in Example 15, the title compound was prepared.

¹H NMR δ (CDCl₃, 250 MHz) 7.79-7.61 (m, 2 H), 7.32-7.14 (m, 2 H), 7.00 (dd, 1 H, J = 1.5 and 8.8 Hz), 6.89-6.67 (m, 1 H), 4.75 (appq, 1 H, J = 8.2 and 16.2 Hz), 4.57 (appt, 1 H, J = 8.6 and 17.4 Hz), 4.49-3.71 (m, 4 H), 3.85 (s, 3 H), 1.92-1.52 (m, 3 H), 1.08-0.78 (m, 6 H).

10

MS calcd for (C₂₇H₃₄N₂O₅S - H)⁺: 403. Found: 403.


Examples 79-93

15

By analogous procedures to those described in Example 15, using the appropriate amino acid and acid or acid chloride reagents consistent with the final products, the compounds of Table 2 were also prepared. ¹H NMR spectra and/or mass spectra were consistent with the structures in Table 2.

20

Table 2



Example	R ³	R ⁴	synthesis method
79		phenyl	SPS
80	"	3-chlorophenyl	SPS
81	"	2-phenylethen-1-yl	SPS
82	"	3-fluorophenyl	SPS
83	"	3-hydroxyphenyl	SPS
84	"	4-methylphenyl	SPS
85	"	4-isopropylphenyl	SPS
86	"	4-trifluoromethylphenyl	SPS
87	"	4-methylthiophenyl	SPS
88	"	4-(benzylsulphonylamino)phenyl	SPS
89	"	4-(diethylaminosulphonyl)phenyl	SPS
90	"	4-(acetylamino)phenyl	SPS
91	"	4-benzoylphenyl	SPS
92	"	4-acetylphenyl	SPS
93	"	4-(4-oxopent-1-yl)phenyl	SPS

5

Example 94Preparation of 4-S-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-one

10

(a) *trans*-4-S-Amino-3-R-hydroxytetrahydrofuran hydrochloride

A mixture of *trans*-4-S-azido-3-R-hydroxytetrahydrofuran (ref L.E. Martinez, J.L. Leighton, D.E. Carsten and E.N. Jacobsen, *J. Amer. Chem. Soc.*, 1995, **117**, 5897) (10 g, 77 mmol) and 10% palladium on charcoal (1 g) in ethanol (150 ml) was stirred under an atmosphere of hydrogen (50 psi) for 12 h. The mixture was filtered and treated with 100 ml of ethanolic HCl to afford, after evaporation under reduced pressure, the title compound as a brown solid, 10.5 g, 97% yield. m.p. 132 °C

¹H NMR δ (D₂O) 4.54-4.52 (m, 1 H), 4.24-4.13 (m, 2 H), 3.98-3.61 (m, 3 H).

15

(b) *trans*-4-*S*-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran

- Trimethylacetyl chloride (5.4 ml, 44 mmol) was added to a stirred solution of N-Cbz-L-leucine (12.7 g, 48 mmol) and triethylamine (14 ml, 52 mmol) in THF (200 ml). After 1 h, *trans*-4-*S*-amino-3-*R*-hydroxytetrahydrofuran.HCl (5.58 g, 40 mmol) was added and the mixture was allowed to stir at reflux for 16 h. The reaction mixture was filtered and evaporated under reduced pressure. Flash column chromatography (80% ethyl acetate-hexane) afforded the title compound as a white foam, 10.6 g, 76% yield.
- ¹H NMR δ (CDCl₃) 7.76 (d, 1 H, J = 5.3 Hz), 7.33-7.20 (m, 5 H), 6.43 (d, 1 H, J = 8.9 Hz), 5.01 (appd, 2 H, J = 3.0 Hz), 4.60-3.65 (m, 6 H), 1.61-1.42 (m, 3 H), 0.93-0.88 (m, 6 H)

(c) *trans*-4-*S*-Amino-N-(*S*-leucine)-3-*R*-hydroxytetrahydrofuran hydrochloride

- A mixture of *trans*-4-*S*-Amino-N-[(carbobenzyloxy)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran (2.0 g, 5.7 mmol and 10% palladium on charcoal (500 mg) in ethanol (100 ml) was stirred under an atmosphere of hydrogen (50 psi) for 12 hours. The reaction mixture was then filtered diluted with ethereal HCl (100 ml, 1 molar) and evaporated under reduced pressure to afford the title compound as a white foam, 1.5 g, 100% yield.
- ¹H NMR δ (D₂O) 4.20-4.05 (m, 2 H), 4.06-4.00 (m, 1 H), 3.90-3.83 (m, 2 H), 3.68-3.52 (m, 2 H), 1.65-1.47 (m, 3 H), 0.86-0.78 (m, 6 H).

(d) *trans*-4-*S*-Amino-N-[(2-benzo(b)thiophenecarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran

- N,N-Diisopropylethylamine (0.4 ml, 2.0 mmol) was added to a stirred solution of *trans*-4-*S*-amino-N-(*S*-leucine)-3-*R*-hydroxytetrahydrofuran.HCl salt (380 mg, 1.1 mmol) in dichloromethane (10 ml). After 5 minutes benzo[b]thiophene-2-carbonyl chloride (196 mg, 1.0 mmol) was added and the mixture was allowed to stir for 1 h then evaporated under reduced pressure. Flash column chromatography (40% acetone-hexane) afforded the title compound as a white foam, 271 mg, 75% yield.
- ¹H NMR δ (d, DMSO, 400 MHz) 8.72 (d, 1 H, J = 8.5 Hz), 8.25 (s, 1 H), 8.04 (d, 1 H, J = 8.5 Hz), 7.95 (d, 1 H, J = 8.5 Hz), 7.46-7.43 (m, 2 H), 5.23 (d, 1 H, J = 3.94 Hz), 4.54-4.47 (m, 1 H), 4.03-4.00 (m, 2 H), 3.90 (dd, 1 H, J = 5.4 and 8.9 Hz), 3.82 (dd, 1 H, J =

4.4 and 9.3 Hz). 3.53- 3.48 (m, 2 H), 1.72-1.66 (m, 2 H), 1.52-1.48 (m, 1 H), 0.91 (d, 3 H, J = 6.4 Hz), 0.88 (d, 3 H, J = 6.4 Hz).

MS calcd for $(C_{18}H_{28}N_2O_5 + H)^+$: 365. Found: 365.

5 (e) 4-S-Amino-N-[(2-benzo(b)thiophenecarbonyl)-S-leucine]-tetrahydrofuran-3-one

Dess-Martin periodinane (200 mg, 0.5 mmol) was added to a stirred solution of *trans*-4-S-Amino-N-[(2-benzo(b)thiophenecarbonyl)-S-leucine-3-R-hydroxytetrahydrofuran (150 mg, 0.40 mmol) in dichloromethane (10 ml). After 1 h, ether (20 ml) was added followed by sodium thiosulfate (1 g). After an additional 15 mins the reaction was washed with saturated sodium hydrogen carbonate, brine and dried. Evaporation of the solvent gave the title compound as a white foam, 147 mg, 100% yield.

1H NMR δ (d_6 DMSO, 400 MHz) 8.82 (d, 1 H, J = 4.0 Hz), 8.46 (d, 1 H, J = 4.0 Hz), 8.28 (s, 1H), 8.05 (d, 1 H, J = 4.0 Hz), 7.98 (d, 1 H, J = 4.0 Hz), 7.47-7.44 (m, 2 H), 4.54-4.51 (m, 1 H), 4.33-4.18 (m, 3 H), 4.08-3.80 (m, 3 H), 1.74-1.67 (m, 2 H), 1.58-1.56 (m, 1 H), 0.92 (d, 3 H, J = 6.4 Hz), 0.88 (d, 3 H, J = 6.4 Hz).

MS calcd for $(C_{19}H_{22}N_2O_4 - H)^+$: 373. Found: 373.

20

Example 95

Preparation of 4-R-Amino-N-[(2-benzo(b)thiophenecarbonyl)-S-leucine]-tetrahydrofuran-3-one

25

Following the procedures of Example 94(a-e), but starting instead with *trans*-4-R-azido-3-S-hydroxytetrahydrofuran (ref L.E. Martinez, J.L. Leighton, D.E. Carsten and E.N. Jacobsen, *J. Amer. Chem. Soc.*, 1995, 117, 5897) in the method of Example 94(a), the title compound was prepared.

1H NMR δ (d_6 DMSO, 400 MHz) 8.82 (d, 1 H, J = 4.0 Hz), 8.52 (d, 1 H, J = 4.0 Hz), 8.28 (s, 1H), 8.01 (d, 1 H, J = 4.0 Hz), 7.95 (d, 1 H, J = 4.0 Hz), 7.47-7.44 (m, 2 H), 4.54-4.51 (m, 1 H), 4.33-4.18 (m, 3 H), 4.33 (appt, 1 H, J = 8.5 Hz), 4.22 (dd, 1 H, J = 8.7 and 16.0 Hz), 4.07 (appd, 1 H, J = 16.6 Hz), 3.90 (appd, 1 H, J = 16.6 Hz), 3.81 (appt, 1 H, J = 8.5 Hz), 1.74-1.67 (m, 2 H), 1.58-1.56 (m, 1 H), 0.92 (d, 3 H, J = 6.4 Hz), 0.88 (d, 3 H, J = 6.4 Hz).

35

MS calcd for $(C_{19}H_{22}N_2O_4 - H)^+$: 373. Found: 373.

Example 96Preparation of 4-S-Amino-N-[(2-naphthyl)-S-leucine]-tetrahydrofuran-3-one

5 Following the procedures of Example 94, using the appropriate carboxylic acid chloride and the 4-azido-3-hydroxytetrahydrofuran of the required stereochemistry to be consistent with the final product, the following compound was prepared.

¹H NMR δ (d, DMSO, 400 MHz) 8.64 (d, 1 H, J = 3.8 Hz), 8.54 (s, 1 H), 8.42 (d, 1 H, J = 3.8 Hz), 8.03 (d, 1 H, J = 3.8 Hz), 8.02-7.95 (m, 3 H), 7.62-7.58 (m, 2 H), 4.63-4.60
10 (m, 1 H), 4.35-4.27 (m, 2 H), 4.08 (appd, 1 H, J = 16.8 Hz), 3.86 (appd, 1 H, J = 16.8 Hz), 3.82 (appt, 1 H, J = 8.0 Hz), 1.75-1.64 (m, 2 H), 1.62-1.55 (m, 1 H), 0.92 (d, 3 H, J = 6.0 Hz), 0.88 (d, 3 H, J = 6.0 Hz).

MS calcd for (C₂₁H₂₄N₂O₄ - H)⁺: 367. Found: 367.

15

Example 97Preparation of 4-R-N-[(2-naphthyl)-S-leucine]-tetrahydrofuran-3-one

20 Following the procedures of Example 94, using the appropriate carboxylic acid chloride and the 4-azido-3-hydroxytetrahydrofuran of the required stereochemistry to be consistent with the final product, the following compound was prepared.

¹H NMR δ (d, DMSO, 400 MHz) 8.62 (d, 1 H, J = 3.8 Hz), 8.52 (s, 1 H), 8.50 (d, 1 H, J = 3.8 Hz), 8.04 (d, 1 H, J = 3.8 Hz), 8.02-7.95 (m, 3 H), 7.62-7.58 (m, 2 H), 4.63-4.60
(m, 1 H), 4.32 (appt, 1 H, J = 8.8 Hz), 4.23 (ddd, 1 H, J = 8.8 and 16.0 Hz), 4.04 (appd, 1
25 H, J = 16.4 Hz), 3.86 (appd, 1 H, J = 16.4 Hz), 3.81 (appt, 1 H, J = 8.0 Hz), 1.75-1.64 (m, 2 H), 1.62-1.55 (m, 1 H), 0.92 (d, 3 H, J = 6.0 Hz), 0.88 (d, 3 H, J = 6.0 Hz).

MS calcd for (C₂₁H₂₄N₂O₄ - H)⁺: 367. Found: 367.

30

Example 98Preparation of 4-S-Amino-N-[(quinoline-2-carbonyl)-S-leucine]-tetrahydrofuran-3-one

35 Following the procedures of Example 94, using the appropriate carboxylic acid chloride and the 4-azido-3-hydroxytetrahydrofuran of the required stereochemistry to be consistent with the final product, the following compound was prepared.

¹H NMR δ (d, DMSO, 400 MHz) 8.80 (d, 1 H, J = 9.0 Hz), 8.70 (d, 1 H, J = 6.9 Hz), 8.63 (d, 1 H, J = 8.5 Hz), 8.39 (d, 1 H, J = 2.5 Hz), 8.27 (d, 1 H, J = 2.5 Hz), 8.21 (d, 1

H, J = 7.9 Hz), 7.95 (appt, 1 H, J = 8.2 Hz), 7.77 (appt, 1 H, J = 7.1 Hz), 4.75-4.66 (m, 1 H), 4.41-4.26 (m, 2 H), 4.16-3.79 (m, 3 H), 1.82-1.63 (m, 3 H), 0.97-0.88 (m, 6 H).
MS calcd for $(C_{30}H_{29}N_3O_4 + H)^+$: 370. Found: 370.

5

Example 99Preparation of 4-R-Amino-N-[(quinoline-2-carbonyl)-S-leucine]-tetrahydrofuran-3-one

- Following the procedures of Example 94, using the appropriate carboxylic acid chloride and the 4-azido-3-hydroxytetrahydrofuran of the required stereochemistry to be consistent with the final product, the following compound was prepared.
- ¹H NMR δ (d₆ DMSO, 400 MHz) 8.88 (d, 1 H, J = 9.0 Hz), 8.82 (d, 1 H, J = 6.7 Hz), 8.70 (d, 1 H, J = 8.5 Hz), 8.30 (d, 1 H, J = 2.5 Hz), 8.27 (d, 1 H, J = 2.5 Hz), 8.21 (d, 1 H, J = 7.9 Hz), 8.02 (appt, 1 H, J = 7.0 Hz), 7.87 (appt, 1 H, J = 7.0 Hz), 4.82-4.73 (m, 1 H), 4.48-3.88 (m, 5 H), 1.88-1.70 (m, 3 H), 1.05-1.02 (m, 6 H).
- MS calcd for $(C_{20}H_{23}N_3O_4 + H)^+$: 370. Found: 370.

Example 100


- 20 Preparation of 4-S-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-one

- Following the procedures of Example 94, using the appropriate carboxylic acid chloride and the 4-azido-3-hydroxytetrahydrofuran of the required stereochemistry to be consistent with the final product, the following compound was prepared.
- ¹H NMR δ (d₆ DMSO, 400 MHz) 8.73 (d, 1 H, J = 8.4 Hz), 8.52 (d, 1 H, J = 7.0 Hz), 7.65-7.62 (m, 2 H), 7.33 (d, 1 H, J = 2.6 Hz), 7.12 (dd, 1 H, J = 2.6 and 9.2 Hz), 4.64-4.55 (m, 1 H), 4.42-4.30 (m, 2 H), 4.18-3.87 (m, 3 H), 3.86 (s, 3 H), 1.84-1.57 (m, 3 H), 0.98-0.87 (m, 6 H).
- 30 MS calcd for $(C_{20}H_{26}N_2O_6 + H)^+$: 389. Found: 389.

Examples 101-108

- By analogous procedures to those described in Example 15, using the appropriate amino acid and acid or acid chloride reagents consistent with the final products, the compounds of Table 3 were also prepared. ¹H NMR spectra and/or mass spectra were consistent with the structures in Table 3.

Table 3



5


Example	R ³	R''	synthesis method
101		6-hydroxybenzo[b]thiophen-2-yl	SPS
102	"	5-hydroxybenzo[b]thiophen-2-yl	SPS
103	"	4-(3-(hydroxymethyl)phenyl)phenyl	SPS
104	"	3-phenylphenyl	SPS
105	"	4-(3-oxophenyl)phenyl	SPS
106	"	4-(3-(aminosulphonyl)phenyl)phenyl	SPS
107		benzo[b]thiophen-2-yl	SPS
108		benzo[b]thiophen-2-yl	SPS

Examples 109-126

Following the procedures of Example 94, using the appropriate carboxylic acid chloride and the 4-azido-3-hydroxytetrahydrofuran of the required stereochemistry to be consistent with the final products, examples 109-126 were prepared. ¹H NMR spectra and/or mass spectra were consistent with the structures in Table 4.

15

Table 4



Example	R ³	R''	stereochem at 4-position
109		4-(pyrid-3-yl)phenyl	S
110	"	4-(pyrid-2-yl)phenyl	S


111	"	4-acetylphenyl	S
112	"	benzyloxy	S
113	"	3,4-dimethoxyphenyl	S
114	"	benzofuran-2-yl	S
115	"	4-(6-methylpyrid-3-yl)phenyl	S
116	"	5-chlorobenzo[b]thiophen-2-yl	S
117	"	4-(pyrid-4-yl)phenyl	S
118	"	2-chlorophenyl	S
119	"	4-bromophenyl	S
120	"	4-chlorobenzo[b]thiophen-2-yl	S
121	"	4-benzylpiperidin-1-yl	S
122	"	3,4-dichlorophenyl	S
123	"	3,4-dimethoxyphenyl	R
124	"	benzofuran-2-yl	R
125	"	3-chlorophenyl	S
126	"	5-chlorobenzo[b]thiophen-2-yl	R

Examples 127-129

- Using the SPS procedure of Example 15 but substituting 3,3-dimethoxy-4-aminotetrahydropyran for 3,3-dimethoxy-4-aminotetrahydrofuran, and using appropriate amino acid and carboxylic acid reagents consistent with the final products, examples 127-129 were prepared. ¹H NMR spectra and/or mass spectra were consistent with the structures in Table 5.

10

Table 5



Example	R ³	R ⁴	synthesis method
127		4-phenoxyphenyl	SPS
128	"	quinolin-2-yl	SPS
129	"	3,4-dimethoxyphenyl	SPS

Example 130Preparation of 4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrothiophen-3-one

5

(a) *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine methyl ester

Benzo[b]thiophene-2-carbonyl chloride (4.9 g, 25 mmol) was added to a stirred solution of L-leucine methyl ester (4.5 g, 25 mmol) and diisopropylethylamine (9 ml, 51 mmol) in DCM (200 ml). After stirring for 2 hours at room temperature, the mixture was poured into water and washed with brine and dried (MgSO₄). Evaporation under reduced pressure afforded the title compound as a white solid, 7.6 g, 100% yield.

¹H NMR δ (CDCl₃, 250 MHz) 7.88-7.78 (m, 3 H), 7.44-7.38 (m, 2 H), 6.53 (d, 1 H, J = 7.6 Hz), 4.91-4.82 (m, 1 H), 3.78 (s, 3 H), 1.82-1.60 (m, 3 H), 1.00 (app t, 6 H, J = 5.9 Hz).

15

(b) *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine

Lithium hydroxide (1.41 g, 59 mmol) was added in one portion to a stirred solution of *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine methyl ester (8.99 g, 29.4 mmol) in THF/H₂O (1/1, 300 ml). After stirring for 12 hours at room temperature, the mixture was poured into water and acidified to pH 1 with cHCl and extracted with Et₂O (x2). Evaporation under reduced pressure afforded the title compound as a yellow solid, 6.1 g, 71% yield.

¹H NMR δ (d₆ DMSO, 250 MHz) 8.78 (d, 1 H, J = 9.1 Hz), 8.12 (s, 1H), 7.92-7.83 (m 2 H), 7.36-7.30 (m, 2 H), 4.37-4.28 (m, 1 H), 1.88-1.47 (m, 3 H), 0.82 (d, 3 H, J = 6.0 Hz), 0.78 (d, 3 H, J = 6.0 Hz).

25

(c) *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine-*S*-(methoxycarbonylmethyl)-L,D-cysteine ethyl ester

30

Iso-Propyl chloroformate (5.9ml, 1.0M in toluene) was added to a stirred solution of *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine (1.65g, 5.7 mmol) and triethylamine (1.6 ml, 11.8 mmol) in DCM (20 ml). After 1 hour at room temperature L-cysteine ethyl ester was added followed by triethylamine (1.6 ml, 11.8 mmol) and the mixture was allowed to stir for a further 6 hours. Methyl bromoacetate (0.6 ml, 6.3 mmol) was added and the mixture was left for a further 0.5 hours before being poured

35

into water and extracted with EtOAc(x3). The combined organic layers were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Flash column chromatography (40% hexane-Et₂O) afforded the title compound as a white solid, 1.6g, 70 %.

- 5 ¹H NMR δ (CDCl₃, 250 MHz) 7.84-7.77 (m, 3 H), 7.66 (d, 0.5 H, J = 8.4 Hz), 7.52 (d, 0.5 H, J = 8.4 Hz), 7.40-7.35 (m, 2 H), 7.24 (d, 0.5 H, J = 8.4 Hz), 7.16 (d, 0.5 H, J = 8.4 Hz), 4.88-4.82 (m, 2 H), 4.23-4.15 (m, 2 H), 3.70 (s, 1.5 H), 3.69 (s, 1. H), 3.36-3.22 (m, 2 H), 3.15-3.08 (m, 2 H), 1.82-1.75 (m, 3 H), 1.28 (t, 1.5 H, J = 4.5 Hz), 1.23 (t, 1.5 H, J = 4.5 Hz), 0.98-0.95 (m, 6 H).

10

(d) 4-(*R,S*)-Amino-N-[(benzo[b]thiophene-2-ylcarbonyl)-*S*-leucine]-tetrahydrothiophene-3-one

- A solution of freshly prepared sodium methoxide (from 75mg of sodium and 2 ml of methanol) was added to *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine-S-
15 (carbomethoxymethyl)-L,D-cysteine ethyl ester (1.5 g, 3.2 mmol) in methanol (2 ml) at room temperature. After 1 hour Et₂O (80 ml) was added, the solution was cooled to 0 °C and the resulting white solid was filtered off. The solid was dissolved in 15ml of glacial acetic acid, 10ml of cHCl and 15 ml of H₂O. This mixture was stirred at reflux for 0.5
20 hours then cooled to room temperature and extracted with CHCl₃ (x5). The combined organic layers were washed with NaHCO₃, brine, dried (MgSO₄) and evaporated under reduced pressure. Flash column chromatography (40% hexane-Et₂O) afforded the title compound as a white solid, 120 mg, 10%.

- ¹H NMR δ (CDCl₃, 400 MHz) 7.88-7.78 (m, 3 H), 7.58 (d, 0.5 H, J = 6.4 Hz), 7.54 (d, 0.5 H, J = 6.4 Hz), 7.42-7.35 (m, 3 H), 4.85-4.80 (m, 1 H), 4.43-4.34 (m, 1 H), 3.34-3.23 (m, 3 H), 3.02-2.91 (m, 1 H), 1.76-1.74 (m, 3 H), 0.98-0.85 (m, 6 H).
25 MS calcd for (C₁₉H₂₂N₂O₃S + H)⁺: 391. Found: 391.

Example 131

30

Preparation of 4-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3,3-dimethoxytetrahydrofuran, diastereomer 1

- a) 4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3,3-dimethoxytetrahydrofuran
35

4-(*R,S*)-Amino-3,3-dimethoxytetrahydrofuran (2.2g, 15mmol), N-benzyloxycarbonyl-L-leucine (3.98g, 15mmol), EDC (3.17g, 16.5mmol) and hydroxyaminobenzotriazole (0.45g, 3.3mmol)

were stirred together in a mixture of dichloromethane and tetrahydrofuran (1:1, 100ml) at ambient temperature for 12hr. The mixture was added to ethyl acetate (300ml), washed with water (2x100ml), dried (MgSO₄) and evaporated to dryness. Chromatography of the resulting oil on silica gel (ethyl acetate / hexane 1:3 to 1:1 gradient) gave the title compound as an oil

5 (5.27g, 89%).

¹H NMR δ (CDCl₃) 7.34 (s, 5H), 6.55 (d, 1H), 5.11 (m, 3H), 4.34 (q, 1H, J = 6.7Hz), 4.20 (m, 2H), 3.77 (s, 2H), 3.51 (m, 1H), 3.26 (d, 3H, J = 2.3Hz), 3.20 (d, 3H, J = 4.7Hz), 1.54 (m, 3H), 0.95 (s, 3H), 0.93 (s, 3H).

10 b) Separate diastereomers of 4-Amino-N-(*S*-leucine)-3,3-dimethoxytetrahydrofuran

4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3,3-dimethoxytetrahydrofuran (5.27g, 13.4mmol) was subjected to hydrogenation at 50psi in methanol containing 10% palladium on carbon. After 3.5 hr, the mixture was filtered through celite and the solvent was removed under reduced pressure. Chromatography of the resulting oil (3.5g) on silica gel (CH₂Cl₂ containing

15 MeOH, 0-4% gradient) separated the title compound into the two pure single diastereomers: diastereomer 1 (faster running) (0.63g). ¹H NMR δ (CDCl₃) 7.76 (d, 1H, J = 7.3Hz), 4.36 (dd, 1H, J = 6.3, 7.2Hz), 4.21 (dd, 1H, J = 6.4, 9.0Hz), 3.80 (s, 2H), 3.57 (dd, 1H, J = 5.3, 9.1Hz), 3.41 (dd, 1H, J = 3.8, 10.1Hz), 3.30 (s, 3H), 3.24 (s, 3H), 1.73 (m, 2H), 1.35 (m, 1H), 0.96

20 (d, 3H, J = 6.7Hz), 0.94 (d, 3H, J = 6.6Hz)

diastereomer 2 (slower running) (0.91g) ¹H NMR δ (CDCl₃) 7.77 (d, 1H, J = 7.1Hz), 4.36 (dd, 1H, J = 6.4, 12.9Hz), 4.21 (dd, 1H, J = 6.5, 9.1Hz), 3.80 (s, 2H), 3.55 (dd, 1H, J = 5.3, 9.0), 3.38 (dd, 1H, J = 4.0, 10.0), 3.30 (s, 3H), 3.25 (s, 3H), 1.72 (m, 2H), 1.38 (m, 1H), 0.96 (d, 3H, J = 6.7Hz), 0.94 (d, 3H, J = 6.8Hz).

25


c) 4-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3,3-dimethoxytetrahydrofuran, diastereomer 1

Saturated sodium bicarbonate solution (5ml) was added to a solution of 4-amino-N-(*S*-leucine)-3,3-dimethoxytetrahydrofuran, diastereomer 1 (0.13g, 0.5mmol) in 1,4-dioxane (5ml) followed by benzo[b]thiophen-2-ylcarbonyl chloride (0.39g, 2mmol). After 45 minutes, the mixture was added to ethyl acetate, washed with water, sodium bicarbonate solution and water. The organic solution was then dried (MgSO₄) and evaporated to give a white solid (0.47g). Chromatography

35 on silica gel (CH₂Cl₂ containing MeOH 0-5% gradient) gave the title compound (0.18g, 89%) as a white solid. ¹H NMR δ (CDCl₃) 7.88 (s, 1H), 7.74 (m, 3H), 7.34 (m, 2H), 7.18 (d, 1H, J = 7.1Hz), 4.82 (m, 1H), 4.38 (dd, 1H, J = 6.4, 12.3Hz), 4.21 (dd, 1H, J = 6.5, 9.2Hz), 3.78 (ABq, 2H, J = 3.8, 9.8Hz), 3.62 (dd, 1H, J = 5.2, 9.1Hz), 1.78 (m, 3H), 0.99 (s, 3H), 0.96 (s, 3H).

Examples 132-142

- By analogous procedures to those described in example 131, using the appropriate diastereomer of 4-amino-N-(S-leucine)-3,3-dimethoxytetrahydrofuran and the carboxylic acid halide reagents consistent with the final products, the compounds of Table 6 were also prepared. ¹H NMR and mass spectra were consistent with the structures in Table 6.

Table 6

Example	R ³	R''	R ^a	R ^{a'}	diastereomer
132		benzo[b]thiophen-2-yl	Me	Me	2
133	"	indol-5-yl	"	"	1
134	"	"	"	"	2
135	"	quinolin-2-yl	"	"	1
136	"	"	"	"	2
137	"	3-bromophenyl	"	"	1
138	"	"	"	"	2
139	"	4-phenoxyphenyl	"	"	1
140	"	"	"	"	2
141	"	indole-6-yl	"	"	2
142	"	benzimidazol-5-yl			1

Example 143

Preparation of 4-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-S-leucine]-3,3-dimethoxytetrahydropyran diastereomer 1

5

a) 4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-S-leucine]-3,3-dimethoxytetrahydropyran

4-(*R,S*)-amino-N-[(benzyloxycarbonyl)-S-leucine]tetrahydropyran-3-one (example 67c, 3.1g, 8.6mmol) was heated at reflux for 12hr with trimethylorthoformate (2.8ml) and *p*-toluenesulphonic acid (0.080g) in methanol (50ml). After removing the solvent under reduced pressure, the resulting oil was chromatographed on silica gel (ethyl acetate / hexane gradient) to give the title compound as a white solid (2.89g, 80%)

¹H NMR δ (CDCl₃) 7.29 (s, 5H), 6.99 (m, 1H), 6.15 (m, 1H), 5.07 (s, 2H), 4.28 (m, 1H), 4.18(m, 1H), 3.52 (m 4H), 3.19 (s, 3H), 3.14 (s, 3H), 1.85 (m, 1H), 1.63 (m, 4H), 0.93 (s, 6H).

15

b) Separate diastereomers of 4-Amino-N-(S-leucine)-3,3-dimethoxytetrahydropyran

4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-S-leucine]-3,3-dimethoxytetrahydropyran was hydrogenated as in example 131b to give, after chromatography, the pure single diastereomers of the title compound:

diastereomer 1: ¹H NMR δ (CDCl₃) 7.75 (d, 1H, J = 8.1Hz), 4.17 (m, 1H), 3.58 (m, 3H), 3.41 (dd, 1H, J = 3.5, 9.7Hz), 3.28 (s, 3H), 3.24 (s, 3H), 1.92 (m, 1H), 1.71 (m, 4H), 1.30 (m, 1H), 0.97 (d, 3H, J = 6.4Hz), 0.94 (d, 3H, J = 6.3Hz).

diastereomer 2: ¹H NMR δ (CDCl₃) 7.8 (s, 1H), 4.16 (m, 1H), 3.59 (m, 3H), 3.49 (m, 1H), 3.28 (s, 3H), 3.24 (s, 3H), 1.92 (m, 1H), 1.72 (m, 4H), 1.41 (m, 1H), 0.97 (d, 3H, J = 6.1Hz), 0.95 (d, 3H, J = 6.0Hz).

25

c) 4-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-S-leucine]-3,3-dimethoxytetrahydropyran, diastereomer 1

30

4-Amino-N-(S-leucine)-3,3-dimethoxytetrahydropyran diastereomer 1 was reacted with benzo[b]thiophen-2-ylcarbonyl chloride, as in example 131c, to give the title compound.

¹H NMR δ (CDCl₃) 7.98 (d, 1H, J = 8.3Hz), 7.93 (s, 1H), 7.78 (d, 1H, J = 7.7Hz), 7.71 (d, 1H, J = 8.3Hz), 7.33 (m, 3H), 4.83 (m, 1H), 4.19 (m, 1H), 3.69 (m, 1H), 3.57 (m, 2H), 3.40 (d, 1H, J = 13Hz), 3.23 (s, 3H), 3.13 (s, 3H), 1.85 (m, 5H), 0.98 (d, 3H, J = 4.8Hz), 0.96 (d, 3H, J = 5.2Hz).

35

Example 144Preparation of 4-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-S-leucine]-3,3-
dimethoxytetrahydropyran diastereomer 2

5


Using diastereomer 2 of 4-amino-N-(S-leucine)-3,3-dimethoxytetrahydropyran in the procedure of example 143c, the title compound was prepared.

10 The above specification and Examples fully disclose how to make and use the compounds of the present invention. However, the present invention is not limited to the particular embodiments described hereinabove, but includes all modifications thereof within the scope of the following claims. The various references to journals, patents and other publications which are cited herein comprise the state of the art and are incorporated herein by reference as though fully set forth.

15

What is claimed is:

1. A compound according to formula (I):



wherein:

R^1 is R'' , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

R^2 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- 10 R^3 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

R^4 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

each R^5 independently is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- 15 R^6 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, $Het-C_{0-6}$ alkyl;

R' is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

R'' is C_{1-6} alkyl, $Ar-C_{0-6}$ alkyl, $Het-C_{0-6}$ alkyl, $Ar-C_{2-6}$ alkenyl; or $Het-C_{2-6}$ alkenyl;

- 20 X is O or S;

n is 1, 2, or 3;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 wherein R^2 and R^4 are each H.

25


3. A compound according to claim 1 wherein R^3 is C_{1-6} alkyl or C_{2-6} alkenyl.

4. A compound according to claim 3 wherein R^3 is i-butyl.

30

5. A compound according to claim 1 wherein R^1 is $R''OC(O)$, $R''SO_2$ or $R''C(O)$, in which R'' is $Ar-C_{0-6}$ alkyl or $Het-C_{0-6}$ alkyl.

6. A compound according to claim 5 wherein the Rⁿ group is



in which B₂ is OH, CN, OCF₃, OC₁₋₆alkyl, OAr, SO₂C₁₋₆alkyl, C₁₋₆alkyl or halo.

5

7. A compound according to claim 1 wherein n is 1 or 2.

8. A compound according to claim 7 wherein n is 1.


10

9. A compound according to claim 1 wherein X is O.


10. A compound according to claim 1 wherein each R⁵ is H.

11. A compound according to claim 1 of the formula (IIa):

15




12. A compound according to claim 1 of the formula (IIb):



(IIb).

13. A compound according to claim 1 of the formula (IIc):



(IIc).

14. A compound according to claim 1 which is:

- 4-(*R,S*)-Amino-N-[(3,4-methylenedioxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(3,4-dichlorobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(2-quinolinecarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(8-quinolinecarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-2,2-dibenzyl-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(benzo[*b*]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(indole-6-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(benzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(5-aminobenzo[*b*]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(5-chlorobenzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(3-bromobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(4-bromobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(5-chlorobenzo[*b*]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(4-fluorobenzo[*b*]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;

- 4-(*R,S*)-Amino-N-[(4-phenoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(4-phenylbenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(6-trifluoromethylbenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 5 4-(*R,S*)-Amino-N-[(4-ethylbenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(4-(tert-butyl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(5-methoxybenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(4-nitrobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 10 4-(*R,S*)-Amino-N-[(6-bromobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(5-bromobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 15 4-(*R,S*)-Amino-N-[(6-methoxybenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(benzo(b)thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*R*-Amino-N-[(benzo(b)thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-
- 20 one;
 4-*S*-Amino-N-[(2-naphthoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*R*-Amino-N-[(2-naphthoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(quinoline-2-carbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*R*-Amino-N-[(quinoline-2-carbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 25 4-*S*-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-pyrid-3-yl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-pyrid-2-yl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 30 4-*S*-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(benzofuran-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(4-[6-methylpyrid-3-yl]benzoyl)-*S*-leucine]-tetrahydrofuran-3-
- one;
 4-*S*-Amino-N-[(5-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-tetrahydrofuran-3-one;
- 35 4-*S*-Amino-N-[(4-pyrid-4-yl)benzoyl)-*S*-leucine]-tetrahydrofuran-3-one;
 4-*S*-Amino-N-[(2-chlorobenzoyl)-*S*-leucine]-tetrahydrofuran-3-one;

- 4-S-Amino-N-[(4-bromobenzoyl)-S-leucine]-tetrahydrofuran-3-one;
 4-S-Amino-N-[(4-chlorobenzo[b]thiophen-2-ylcarbonyl)-S-leucine]-
 tetrahydrofuran-3-one;
 4-S-Amino-N-[(4-benzylpiperidin-1-ylcarbonyl)-S-leucine]-tetrahydrofuran-3-
 5 one;
 4-S-Amino-N-[(3,4-dichlorobenzoyl)-S-leucine]-tetrahydrofuran-3-one;
 4-S-Amino-N-[(3-chlorobenzoyl)-S-leucine]-tetrahydrofuran-3-one;
 4-(*R,S*)-Amino-N-[(3,4-dimethoxybenzoyl)-S-leucine]-tetrahydropyran-3-one;
 4-(*R,S*)-Amino-N-[(4-phenoxybenzoyl)-S-leucine]-tetrahydropyran-3-one;
 10 4-(*R,S*)-Amino-N-[(quinolin-2-ylcarbonyl)-S-leucine]-tetrahydropyran-3-one;
 4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-S-leucine]-tetrahydropyran-3-one;
 4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-S-leucine]-
 tetrahydropyran-3-one; or
 4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-S-leucine]-
 15 tetrahydrothiophen-3-one;
 or a pharmaceutically acceptable salt thereof.

15. A pharmaceutical composition comprising a compound according to
 any one of claims 1-14 and a pharmaceutically acceptable carrier.

20

16. A method of inhibiting a cysteine protease which comprises
 administering to a patient in need thereof an effective amount of a compound according
 to claim 1.

25

17. A method according to claim 16 wherein the cysteine protease is
 cathepsin K.

30

18. A method of inhibiting bone loss which comprises administering to a
 patient in need thereof an effective amount of a compound according to claim 1.

35


19. A method of treating osteoporosis which comprises administering to a
 patient in need thereof an effective amount of a compound according to claim 1.

40

20. A method of treating gingival or periodontal disease which comprises
 administering to a patient in need thereof an effective amount of a compound according to
 claim 1.

21. A method of treating a disease characterized by excessive cartilage or matrix degradation which comprises administering to a patient in need thereof an effective amount of a compound according to claim 1.
- 5
22. A method according to claim 21 wherein said disease is osteoarthritis or rheumatoid arthritis.
23. A compound according to any one of claims 1 to 14 for use as a
- 10 medicament.
24. The use of a compound of the formula (I) as defined in claim 1 in the manufacture of a medicament for the treatment of diseases in which inhibition of a cysteine protease is a factor.
- 15
25. The use of a compound according to claim 24 wherein the cysteine protease is cathepsin K.
26. The use of a compound of the formula (I) as defined in claim 1 in the
- 20 manufacture of a medicament for the inhibition of bone loss.
27. The use of a compound of the formula (I) as defined in claim 1 in the manufacture of a medicament for the treatment of osteoporosis.
- 25
28. The use of a compound of the formula (I) as defined in claim 1 in the manufacture of a medicament for the treatment of gingival or periodontal disease.
29. The use of a compound of the formula (I) as defined in claim 1 in the
- 30 manufacture of a medicament for the treatment of diseases characterized by excessive cartilage or matrix degradation.
31. The use of a compound according to claim 29 wherein the disease
- characterized by excessive cartilage or matrix degradation is osteoarthritis or
- 35 rheumatoid arthritis.
31. A process for preparing a compound of the formula (I) as defined in claim 1, which process comprises:

- (i) reacting a compound of the formula (III):




5

(III)

wherein R^1 , R^2 , R^3 , R^4 , R^5 and n are as defined in formula (I) of claim 1, with any reactive functional groups protected, with an oxidizing agent; or


- 10 (ii) decarboxylating a compound of the formula (IV):



(IV)

- 15 wherein R^1 , R^2 , R^3 , R^4 , R^5 and n are as defined in formula (I) of claim 1, with any reactive functional groups protected; or

- (iii) reacting a compound of the formula (V):



20


(V)

wherein R^1 , R^3 , R^4 , R^5 and n are as defined in formula (I) of claim 1, with any reactive functional groups protected, with an acid;

25

and thereafter removing any protecting groups and optionally forming a pharmaceutically acceptable salt.

32. A compound according to formula (III):



- 5 wherein:

R^1 is R'' , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

R^2 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- 10 R^3 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

R^4 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

each R^5 independently is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- 15 R^6 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, $Het-C_{0-6}$ alkyl;


R' is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

R'' is C_{1-6} alkyl, $Ar-C_{0-6}$ alkyl, $Het-C_{0-6}$ alkyl, $Ar-C_{2-6}$ alkenyl; or $Het-C_{2-6}$ alkenyl; and

n is 1, 2 or 3;

- 20 or a pharmaceutically acceptable salt thereof.

33. A compound according to formula (IV):



- 25 wherein:


R^1 is R'' , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

R^2 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- 30 R^3 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- R^4 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
 each R^5 independently is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or
 Het-C₀₋₆alkyl;
 R^6 is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl-C₀₋₆alkyl, Ar-C₀₋₆alkyl,
 5 Het-C₀₋₆alkyl;
 R^7 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
 R^8 is C₁₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, Ar-C₂₋₆alkenyl; or
 Het-C₂₋₆alkenyl; and
 n is 1, 2 or 3;
 10 or a pharmaceutically acceptable salt thereof.

34. A compound according to formula (V):




- 15 wherein:
 R^1 is R^{''}, R^{''}C(O), R^{''}C(S), R^{''}SO₂, R^{''}OC(O), R^{''}R^{''}NC(O), or
 R^{''}OC(O)NR^{''}CH(R⁶)C(O);
 R^3 is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆cycloalkyl-C₀₋₆alkyl,
 Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
 20 R^4 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
 each R^5 independently is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or
 Het-C₀₋₆alkyl;
 R^6 is H, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl-C₀₋₆alkyl, Ar-C₀₋₆alkyl,
 Het-C₀₋₆alkyl;
 25 R^7 is H, C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₀₋₆alkyl, or Het-C₀₋₆alkyl;
 R^8 is C₁₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, Ar-C₂₋₆alkenyl; or
 Het-C₂₋₆alkenyl;
 X is O or S; and
 n is 1, 2 or 3;
 30 or a pharmaceutically acceptable salt thereof.

35. A compound selected from the group consisting of:
trans-4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3-
 hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-[(*tert*-butoxycarbonyl)-*S*-leucine]-3-
 5 hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-(*S*-leucine)-3-hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-[(3,4-methylenedioxybenzoyl)-*S*-leucine]-3-
 hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-[(3,4-dichlorobenzoyl)-*S*-leucine]-3-
 10 hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-[(2-quinolinecarbonyl)-*S*-leucine]-3-
 hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3-
 hydroxytetrahydropyran;
 15 *trans*-4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-
 hydroxytetrahydropyran;
trans-4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-
 hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-[(indole-6-ylcarbonyl)-*S*-leucine]-3-
 20 hydroxytetrahydrofuran;
trans-4-(*R,S*)-Amino-N-[(5-aminobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-
 hydroxytetrahydrofuran
trans-4-Amino-3-hydroxytetrahydrofuran;
trans-3-Hydroxy-4-benzyloxycarbonylamino-tetrahydrofuran;
 25 4-Benzyloxycarbonylamino-tetrahydrofuran-3-one;
 3,3-Dimethoxy-4-benzyloxycarbonylamino-tetrahydrofuran;
 3,3-Dimethoxy-4-amino-tetrahydrofuran
trans-4-*S*-Amino-3-*R*-hydroxytetrahydrofuran;
trans-4-*S*-Amino-N-[(benzyloxycarbonyl)-*S*-leucine]-3-*R*-
 30 hydroxytetrahydrofuran;
trans-4-*S*-Amino-N-(*S*-leucine)-3-*R*-hydroxytetrahydrofuran;
trans-4-*S*-Amino-N-[(3,4-dichlorobenzoyl)-*S*-leucine]-3-*R*-
 hydroxytetrahydrofuran;
trans-4-*S*-Amino-N-[(2-quinolinecarbonyl)-*S*-leucine]-3-*R*-
 35 hydroxytetrahydrofuran;
trans-4-*S*-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-*R*-
 hydroxytetrahydrofuran;

- trans*-4-*S*-Amino-N-[(benzofuran-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(2-naphthoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(5-methoxybenzofuran-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- 5 *trans*-4-*S*-Amino-N-[(5-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(4-chlorobenzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- 10 *trans*-4-*S*-Amino-N-[(4-bromobenzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(4-(pyrid-2-yl)benzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-*S*-Amino-N-[(4-(pyrid-3-yl)benzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- 15 *trans*-4-*S*-Amino-N-[(3,4-dimethoxybenzoyl)-*S*-leucine]-3-*R*-hydroxytetrahydrofuran;
- trans*-4-Amino-3-hydroxytetrahydropyran;
- trans*-4-(*R,S*)-Amino-N-[(benzoyloxycarbonyl)-*S*-leucine]-3-hydroxytetrahydropyran;
- 20 *trans*-4-(*R,S*)-Amino-N-(*S*-leucine)-3-hydroxytetrahydropyran;
- trans*-4-(*R,S*)-Amino-N-[(benzo[b]thiophen-2-ylcarbonyl)-*S*-leucine]-3-hydroxytetrahydropyran;
- N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine methyl ester;
- N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine;
- 25 *N*-benzo[b]thiophene-2-ylcarbonyl-L-leucine-*S*-(methoxycarbonylmethyl)-L,D-cysteine ethyl ester; or
- 2-Methoxycarbonyl-4-(*R,S*)-Amino-N-[(benzo[b]thiophene-2-ylcarbonyl)-*S*-leucine]-tetrahydrothiophene-3-one;
- or salts thereof.

30

36. A compound according to formula (VI):



- 5 wherein:

R^1 is R , $R''C(O)$, $R''C(S)$, $R''SO_2$, $R''OC(O)$, $R''R'NC(O)$, or $R''OC(O)NR'CH(R^6)C(O)$;

R^2 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- 10 R^3 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

R^4 is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

each R^5 independently is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

- 15 R^6 is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} cycloalkyl- C_{0-6} alkyl, $Ar-C_{0-6}$ alkyl, $Het-C_{0-6}$ alkyl;

R' is H, C_{1-6} alkyl, C_{2-6} alkenyl, $Ar-C_{0-6}$ alkyl, or $Het-C_{0-6}$ alkyl;

R'' is C_{1-6} alkyl, $Ar-C_{0-6}$ alkyl, $Het-C_{0-6}$ alkyl, $Ar-C_{2-6}$ alkenyl; or $Het-C_{2-6}$ alkenyl;

X is O or S;

- 20 n is 1, 2 or 3; and

R^A and $R^{A'}$ independently are H or C_{1-2} alkyl, with the proviso that when one of R^A or $R^{A'}$ is H, the other is C_{1-2} alkyl; or together are $(CH_2)_{2-3}$ forming a 5- or 6-membered ring;

or a pharmaceutically acceptable salt thereof.

25

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/03200

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : Please See Extra Sheet.		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched none		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, STN, WPI, CAPLUS, REGISTRY, MEDLINE, BIOSIS		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	WO 97/32846 A1 (PHARMACIA & UPJOHN COMPANY) 12 September 1997, see entire document, especially pages 2-7.	1-36
Y	GB 2 282 598 A (MERCK & CO., INC.) 12 April 1995, see entire document, especially pages 9-13.	1-36
A,P	WO 98/04539 A1 (MITSUBISHI CHEMICAL CORPORATION) 02 May 1998, see entire document, especially the abstract and pages 12-44.	1-36
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 17 JULY 1998		Date of mailing of the international search report 08 SEP 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer <i>Susan Hanley</i> SUSAN HANLEY Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/03200

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12N 9/99, 9/50; A01N 43/02, 43/06, 43/08, 43/16, 43/18; C07D 307/00, 307/48, 309/00, 319/06, 321/00, 333/02, 333/12, 333/22, 335/00, 335/02

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/184, 219; 549/1, 13, 28, 29, 74, 76, 77, 200, 356, 372, 373, 375, 429, 467, 468; 514/430, 432, 438, 447, 448, 449, 451, 461, 472, 473

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

435/184, 219; 549/1, 13, 28, 29, 74, 76, 77, 200, 356, 372, 373, 375, 429, 467, 468; 514/430, 432, 438, 447, 448, 449, 451, 461, 472, 473